

Jla wbgšY Kigwii 19 Riwymii 2011 ZwitL Abn̄Z 239 Zg mfwii Kh̄eei Yx

"T" I cni evi Kj "Y gšYj tqi mPe Rbve gya ūgvqj Knei Gümfvic ZtZi Jla mbqŠY Kigui 239 Zg mfvi MZ  
19 Rbvgui 2011 Zwi tL tej v 12:00 NUKvq gšYj tqi mfvi Kt AbiôZ nq |

*mfig KingUj ibgenV* সদস্যগণ উপস্থিত ছিলেন (জ্যোতির ক্রমানুসারে নয়) :

- 1| W. cÖY tMvciy `E, DcivPih©eVzklL gyRe tgWtKj nekje` yj q, XvKv|

2| tgRi tRbvitij tgvt iweDj tnvtmb, Kbmij tBu wdlRikqib, evsj vt` k AvgW tdtmm tgWtKj mwmfmm|

3| Wt mowm P, ^mvni, Dc-cwi Pyj K (cikimb), ct¶ gnvcwi Pyj K, ^r- Awa`Bi, XvKv|

4| Wt tgvt AvBbjy nK, ct¶ gnvcwi Pyj K, cÖYx mowm Awa`Bi, XvKv|

5| Aa`vcK dwi` v teMg, wKbKij dvgfKij I dvgfKij wR wefim, XvKv nekje` yj q|

6| Aa`vcK tgvt AwRRjy Kvnkvi, tgWtmb wefim, XvKv tgWtKj Ktj R|

7| অধ্যাপক ডাঃ এ কে এম শরিফুল ইসলাম, চর্চ ও ঘোন রোগ বিশেষজ্ঞ, স্যার সলিমুলগ্রাহ মেডিকেল কলেজ, ঢাকা।

8| ডাঃ শামসুন নাহার বেগম, বিভাগীয় প্রধান, গাইনী ও অবস্টেট্রি বিভাগ, স্যার সলিমুলগ্রাহ মেডিকেল কলেজ, ঢাকা।

9| Aa`vcK Wv. tgvt BmgvBj Lvb, cÖZbila, dvgfKij wR wefim, XvKv tgWtKj Ktj R|

10| Wv. Avdmvbw Kvig, ct¶ mfvcwZ, evsj vt` k tgWtKj Gtmwntqkb, XvKv|

11| Wv. tgvtgbjy nK, wmbqi mn mfvcwZ, evsj vt` k JIa lkí migwZ Ges eee`vcbv cwi Pyj K, tRbvitij dvgfimDilUKij m® yj t|

12| nvKig tgvt BDmgd nvi " b fBiqv, BDbvbx wektiA, cÖZbila, evsj vt` k BDbvbx I AvqteP K tevW®

13| tgRi tRbvitij tgvt Avej Kij vq AvRv, gnvcwi Pyj K, JIa cikimb Awa`Bi, XvKv|

*mfvq Avtj vP' ve lq mgn-ibgjeC t*

- 1/ MZ 22-10-2009 Zwi tL AbjōZ JIa mbqšY Kigui 238 Zg mfvi Kvheei Yx bñDZKi Y cñt½ |  
2/ JIa mbqšY Kigui tUKubK'ij mve Kigui MZ 05-05-2010 Zwi tL AbjōZ 56 Zg mfvi mgvwi kmgtni  
wet tq Avtj Pbr I m×všl MōY cñt½ |  
3/ -r- I cwievi Kj 'vY gšyij tqi -yjK bs-Rb -r- -1/JIa-6/2007/329, Zwi L 24-11-2010 tgvZiteK AñZ  
DcKvix cñY kKb i ¶v Ges Añb-ñ mn Abñb tñ vñMi cñKvc eÜKivi j t¶v evsj vñt k ci' PñKrmvq eñeÜZ  
Wbñt KñdbñK RvZxq JIa Drcv`b eÜ Kvievi Ri " ix c' t¶c MōY Kiv cñt½ |  
4/ সভাপতি মহোদয়ের অনমতিক্রমে যে কোন বিষয়।

*mfig DciID<sup>3</sup> relamgr-ve -wi Z Avfj Pbv ceR ibgevZ mowik/mn×všÍMÖY Kiv nBj t*

- 1 | MZ 22-10-2009 Zwitl AbnöZ JIa bgsY KqgjU 238 Zg mfvi Kvh@ei Yx mboZKi Y cñt½

*Aut vPbv t*

*mfvq 238 Zg mfvi Kvh<sup>eei</sup>Yx Dc<sup>-</sup>wcZ nq/ m<sup>i</sup>Pe g<sup>i</sup>nv<sup>i</sup> `q etj b `xN<sup>i</sup>P b ci mfv Ab<sup>i</sup>j<sup>o</sup>Z nBtZtQ/ D<sup>3</sup> mfv Av<sup>i</sup>tv Nb Nb Ab<sup>i</sup>j<sup>o</sup>Z nl qv c<sup>i</sup>qvRb/ JIa w<sup>i</sup>kí mwgwZi w<sup>i</sup>mwbqi mn-mfvciZ Wrt tgwgbj<sup>i</sup> nK etj b `xN<sup>i</sup>eiwZi ci mfv Ab<sup>i</sup>j<sup>o</sup>Z nBtj t`k bZ<sup>i</sup> bZ<sup>i</sup> JIa nBtZ ew<sup>i</sup>AZ nq/*

MZ 22-10-2010 ZwitL AbjöZ 238 mfvi KvheeiYx h\_vh\_fite wjwce× Kiv nBqvtQ ewj qv m`m`MY  
gZcKik Ktib|

im×všít

- (1) *Jla* *nbqšY* *Kiv*<sup>i</sup>*Ui* 238 Zg *mfvi* *Kiv*<sup>ie</sup>*i* *Yx* *nb*<sup>ie</sup>*oZ* *Kiv* *nq* /  
(2) *Avtiv* *Nb* *Nb* *mfvi* *Abötib* *e*<sup>ie</sup>*-v* *MötYi* *Rb*<sup>ie</sup> *gnicwi* *Pvj* *K*, *Jla* *cükimb* *Awa*<sup>ie</sup>*-Bi*<sup>ie</sup>*tK* *Abtiva* *Kiv* *nq* /

2/ *mfvq wbtgæwVZ -vbxq Drcv`b I Avg`vbx Rb` bZb JIa I cPj Z JIta i bZb giv i JIa (mDg`b I tfiUwi bvix) Ges Avg`vbx Rb` tgWtKj WfBtmm Gi tiRt÷kb mæutK®*  
*we -vvi Z Avtj vPbv Kwi qv wbtgæwVZ m×vš/MpxZ nq t*

## 2.1 Proposed Product for Locally Manufacture ( Human)

bs	cÖZKvi tKi bug	JIta i bug I fRtbwK bug	wbt`Rbv	Status (New Molecule/ Existing)	Avt`bKvix cÖE BNF or USFDA Reference	fUKibK`yj mve-KugwJi 56 Zg mfvi m×vš-	mfvi m×vš-
01.	The Ibn Sina Pharmaceuticals Ltd.	a) <b>Tadalafil 20 mg Tablet</b>  Tadalafil INN 20 mg	Treatment of pulmonary arterial hypertension.	New	USFDA	Ace`envi nBtZ cvti weaq mwgmRK tcifvcU wePbv Kwi qv Avte`b -MZ ivLv hvBtZ cvti	Ace`envi nBtZ cvti weaq mwgmRK tcifvcU wePbv Kwi qv Avte`b -MZ ivLv nBj
	b) <b>Milnacipran HCl 25 mg Tablet</b>  Milnacipran Hydrochloride INN 25 mg	It is indicated for the management of fibromyalgia.		New	USFDA	cØqRb bvB weaq Avte`b bvgÄy Kiv hvBtZ cvti	cØqRb bvB weaq Avte`b bvgÄy Kiv nBj
	c) <b>Milnacipran HCl 50 mg Tablet</b>  Milnacipran Hydrochloride INN 50 mg	-do-		New	USFDA	Abfgv`b Kiv hvBtZ cvti	Abfgv`b Kiv nBj
	d) <b>Lacosamide 100 mg Tablet</b>  Lacosamide INN 100 mg	Indicated as adjunctive therapy in the treatment of partial-onset seizures in patients with epilepsy aged 17 years and older and in the treatment of diabetic neuropathic pain.		New	USFDA	cØqRb bvB weaq Avte`b bvgÄy Kiv hvBtZ cvti	cØqRb bvB weaq Avte`b bvgÄy Kiv nBj
	e) <b>Lacosamide 200 mg Tablet</b>  Lacosamide INN 200 mg	-do-		New	USFDA	cØqRb bvB weaq Avte`b bvgÄy Kiv hvBtZ cvti	cØqRb bvB weaq Avte`b bvgÄy Kiv nBj
	f) <b>Saxagliptin 2.5 mg Tablet</b>  Saxagliptin Hydrochloride INN 2.79 mg (eq. to 2.5 mg Saxagliptin)	Indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus.		New	USFDA	Abfgv`b Kiv hvBtZ cvti	Abfgv`b Kiv nBj
	g) <b>Saxagliptin 5 mg Tablet</b>  Saxagliptin Hydrochloride INN 5.58 mg (eq. to Saxagliptin 5 mg)	-do-		New	USFDA	Abfgv`b Kiv hvBtZ cvti	Abfgv`b Kiv nBj

<i>bs</i>	<i>cōZKvičKi bug</i>	<i>Jl̄ai bug I tR̄bniK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigilji 56 Zg mfvi m̄xvišl</i>	<i>mfvi m̄xviš-</i>
	The Ibn Sina Pharmaceuticals Ltd.	h) <b>Telavancin 250 mg/Vial IV Injection</b>  Telavancin Hydrochloride INN 255.308 mg (eq. to Telavancin 250 mg)/Vial	Complicated skin and skin structure infections.	New	USFDA	<i>Abfgv` b Kiv h̄BtZ c̄ti  </i>	<i>Abfgv` b Kiv nBj  </i>
		i) <b>Telavancin 750 mg/Vial IV Injection</b>  Telavancin Hydrochloride INN 765.924 mg (eq. to Telavancin 750 mg)/Vial	Complicated skin and skin structure infections.	New	USFDA	<i>Abfgv` b Kiv h̄BtZ c̄ti  </i>	<i>Abfgv` b Kiv nBj  </i>
		j) <b>Prasugrel 5 mg Tablet</b>  Prasugrel Hydrochloride INN 5.49 mg (Eq. to Prasugrel 5 mg)	Indicated for the prevention of atherothrombotic events in patients with acute coronary syndrome.	New	USFDA	<i>Abfgv` b Kiv h̄BtZ c̄ti  </i>	<i>Abfgv` b Kiv nBj  </i>
		k) <b>Prasugrel 10 mg Tablet</b>  Prasugrel Hydrochloride INN 10.98 mg (Eq. to Prasugrel 10 mg)	-do-	New	USFDA	<i>Abfgv` b Kiv h̄BtZ c̄ti  </i>	<i>Abfgv` b Kiv nBj  </i>
		l) <b>Ganciclovir 0.15% w/w Ophthalmic Gel</b>  Ganciclovir INN 0.150 gm/100 gm	Indicated for the treatment of acute herpetic keratitis (dendritic ulcers).	New	USFDA	<i>Abfgv` b Kiv h̄BtZ c̄ti  </i>	<i>Abfgv` b Kiv nBj  </i>
		m) <b>Bepotastine Besilate 1.5% w/v Ophthalmic Solution</b>  Bepotastine Besilate INN 1.50 gm/100 ml	Indicated in itching associated with signs and symptoms of allergic conjunctivitis.	New	USFDA	<i>Abfgv` b Kiv h̄BtZ c̄ti  </i>	<i>Abfgv` b Kiv nBj  </i>
		n) <b>Cyclobenzaprine Hydrochloride 15 mg Extended Release Capsule</b>  Cyclobenzaprine Hydrochloride Extended release pellets Potency 6% Ph. Gr. 250 mg Contains Cyclobenzaprine Hydrochloride USP 15 mg	Indicated as an adjunct to rest and physical therapy for relief of muscle spasm associated with acute, painful musculoskeletal conditions.	New	USFDA	<i>c̄qyRb b̄B neaq Avt` b b̄gAy Kiv h̄BtZ c̄ti  </i>	<i>c̄qyRb b̄B neaq Avt` b b̄gAy Kiv nBj   </i>

<i>bs</i>	<i>cōZKvi tKi bug</i>	<i>Jl̄tai bug I tR̄tbuiK bug</i>	<i>ib̄t` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt̄e` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK̄yj mve-Kigilji 56 Zg mfvi imxvs̄l</i>	<i>mfvi imxvs̄-</i>
	The Ibn Sina Pharmaceuticals Ltd.	<p>o) <b>Cyclobenzaprine Hydrochloride 30 mg Extended Release Capsule</b>  Cyclobenzaprine Hydrochloride Extended release pellets Potency 12% Ph. Gr. 250 mg contains Cyclobenzaprine Hydrochloride 30 mg</p> <p>p) <b>Rifamixin 200mg Tablet</b>  Rifamixin INN 200mg</p>	-do-	New	USFDA	<i>cōqirB b̄B neaq Avte`b bigĀy Kiv hvBtZ cv̄i /</i>	<i>cōqirB b̄B neaq Avte`b bigĀy Kiv nBj /</i>
02.	Orion Laboratories Ltd.	<p>a) <b>Magaldrate 480 mg + Simethicone 20 mg Chewable Tablet</b>  Magaldrate USP 480 mg + Simethicone USP 20 mg</p> <p>b) <b>Magaldrate 1080 mg + Simethicone 30 mg DS Chewable Tablet</b>  Magaldrate USP 1080 mg + Simethicone USP 30 mg</p> <p>c) <b>Iron Sucrose 200 mg/10 ml Ampoule</b>  Iron Sucrose USP 5.13 gm (eq. to Elemental Iron 200 mg)/10 ml</p> <p>d) <b>Butamirate Citrate 15 mg/10 ml Syrup</b>  Butamirate Citrate BP 150 mg/100 ml</p>	<p>It is indicated for Stomach Ulcer, Ulcer of Duodenum.</p> <p>It is indicated for Stomach Ulcer, Ulcer of Duodenum.</p> <p>Anaemic pregnant women, women of child bearing age specially those with heavy menstrual loss.</p> <p>It is indicated in acute cough of any etiology, pre and post operative cough sedation for surgical procedures and bronchoscopy.</p>	<p>480 mg + 20 mg/5 ml Suspension</p> <p>-do-</p> <p>100 mg/5 ml Injection</p> <p>50 mg Tablet</p>		<i>Abt̄ḡ`b Kiv hvBtZ cv̄i /</i> <i>cōqirB b̄B neaq Avte`b bigĀy Kiv hvBtZ cv̄i /</i> <i>cōqirB b̄B neaq Avte`b bigĀy Kiv hvBtZ cv̄i /</i> <i>USFDA KZR.Abt̄ḡ`Z bq Ges BNF-G Aš-f̄y bq neaq Avte`b bigĀy Kiv hvBtZ cv̄i /</i>	<i>Abt̄ḡ`b Kiv nBj /</i> <i>cōqirB b̄B neaq Avte`b bigĀy Kiv nBj /</i> <i>cōqirB b̄B neaq Avte`b bigĀy Kiv nBj /</i> <i>USFDA KZR.Abt̄ḡ`Z bq Ges BNF-G Aš-f̄y bq neaq Avte`b bigĀy Kiv nBj /</i>

<i>bs</i>	<i>cōZKvi tKi bug</i>	<i>Jl̄tai bug I tR̄tbuiK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigij 56 Zg mfvi imxviši</i>	<i>mfvi imxviš-</i>
03.	Square Pharmaceuticals Ltd., Pabna	<p>a) <b>Naproxen 1000 mg SR Tablet</b> Naproxen Sodium USP 1095.50 mg (eq. to 1000 mg Naproxen USP)</p> <p>b) <b>Cetirizine HCl 5 mg + Pseudoephedrine HCl 120 mg ER Tablet</b> Cetirizine Hydrochloride BP 5 mg + Pseudoephedrine Hydrochloride USP 120 mg</p> <p>c) <b>Light Liquid Paraffin 3% + White Soft Paraffin 8% Lotion</b> Light Liquid Paraffin BP 3 gm + White Soft Paraffin BP 8 gm/100 gm</p> <p>d) <b>Mifepristone 200 mg Tablet</b> Mifepristone INN 200 mg</p> <p>e) <b>Alfacalcidol 0.50 mcg LFHG Capsule</b> Alfacalcidol BP 0.50 mcg</p> <p>f) <b>Alfacalcidol 1 mcg LFHG Capsule</b> Alfacalcidol BP 1 mcg</p> <p>g) <b>Alfacalcidol 2 mcg/ml Injection Ampoule</b> Alfacalcidol BP 2 mcg/ml</p>	Treatment of rheumatoid arthritis, osteoarthritis, ankylosing spondylitis and for the relief of chronic pain.  Indicated for the relief of nasal and non-nasal symptoms associated with seasonal or perennial allergic rhinitis in adults and children of 12 years old.  To treat dry skin conditions.  It is indicated for the medical termination of intrauterine pregnancy through 49 days.  Renal osteodystrophy, Hyperparathyroidism, Neonatal hypocalcaemia, Nutritional and malabsorptive rickets and osteomalacia.	250 mg & 500 mg Tab. & Suppository  New  New  New  250 mg Cap. 1 mcg Tab. 0.500 mcg Tab.	USFDA  USFDA  USFDA  USFDA BNF-57  BNF-57 Page -541 (as Capsule)  BNF-57 Page -542 (as Capsule)  BNF-57 Page -542	<i>Abtgw` Z bq Ges BNF-G Aš-fy bq neaq Avte`b bugÄy Kiv hBtZ cti  </i>  <i>cōqRb bvB neaq Avte`b bugÄy Kiv hBtZ cti  </i>  <i>Abtgw` b Kiv hBtZ cti  </i>  <i>cōqRb bvB neaq Avte`b bugÄy Kiv hBtZ cti  </i>  <i>Abtgw` b Kiv hBtZ cti  </i>	<i>USFDA KZR Abtgw` Z bq Ges BNF-G Aš-fy bq neaq Avte`b bugÄy Kiv nBj  </i>  <i>cōqRb bvB neaq Avte`b bugÄy Kiv nBj  </i>  <i>Abtgw` b Kiv nBj  </i>  <i>cōqRb bvB neaq Avte`b bugÄy Kiv nBj  </i>  <i>Abtgw` b Kiv nBj  </i>  <i>Abtgw` b Kiv nBj  </i>  <i>Abtgw` b Kiv nBj  </i>  <i>Abtgw` b Kiv nBj  </i>

<i>bs</i>	<i>cōZKvi tKi bvg</i>	<i>Jl̄tai bvg I tR̄tbwK bvg</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avtē` bKvix cōĒ BNF or USFDA Reference</i>	<i>tUKibK̄yj mve-KigwJi 56 Zg mfvi wmxvši</i>	<i>mfvi wmxvš-</i>
Square Pharmaceuticals Ltd., Pabna	<p>h) <b>Alfacalcidol 2 mcg/ml Oral Drops</b> Alfacalcidol BP 2 mcg/ml</p> <p>i) <b>Pioglitazone 15 mg + Metformin HCl 1000 mg Extended Release Tablet</b> Pioglitazone Hydrochloride INN 16.530 mg (eq. to Pioglitazone 15 mg) + Metformin Hydrochloride Extended Release Granules 60% w/w Ph. Gr. 1666.667 mg (contains : Metformin Hydrochloride BP 1000 mg)</p> <p>j) <b>Pioglitazone 30 mg + Metformin HCl 1000 mg Extended Release Tablet</b> Pioglitazone Hydrochloride INN 33.060 mg (eq. to Pioglitazone 30 mg) + Metformin Hydrochloride Extended Release Granules 60% w/w Ph. Gr. 1666.667 mg (Contains : Metformin Hydrochloride BP 1000 mg)</p> <p>k) <b>Loperamide HCl 1 mg/5ml Syrup</b> Loperamide Hydrochloride USP 1 mg/5ml</p> <p>l) <b>Racecadotril 100 mg Capsule</b> Racecadotril INN 100 mg</p>	<p>Renal osteodystrophy, Hyperparathyroidism, Neonatal hypocalcaemia, Nutritional and malabsorptive rickets and osteomalacia.</p> <p>Indicated as an adjunct to diet and exercise to improve glycemic control in patients with type 2 diabetes who are already treated with a combination of pioglitazone and metformin.</p> <p>-do-</p> <p>Indicated for the control and symptomatic relief of acute diarrhea and of chronic diarrhea associated with inflammatory bowel disease.</p> <p>Indicated for symptomatic treatment supplementary oral rehydration in acute diarrhea.</p>	<p>250 mg Cap. 1 mcg Tab. 0.500 mcg Tab.</p> <p>Metformin 500 mg + Pioglitazone 15 mg Tablet &amp; Metformin 850 mg + Pioglitazone 15 mg Tablet</p> <p>-do-</p> <p>2 mg Capsule</p>	<p>BNF-57 Page -533</p> <p>USFDA</p> <p>USFDA</p> <p>BNF-57 USFDA</p> <p>New</p>	<p><i>Abtgi` b Kiv hbZ cti /</i></p> <p><i>cōqirb bvB neaq Avte` b bvgÄy Kiv hbZ cti /</i></p> <p><i>cōqirb bvB neaq Avte` b bvgÄy Kiv nbj /</i></p> <p><i>cōqirb bvB neaq Avte` b bvgÄy Kiv hbZ cti /</i></p> <p><i>4(Pri) e0ti i ibtP ikit`i t¶t e eni SAKY neaq Avte` b bvgÄy Kiv hbZ cti /</i></p> <p><i>4(Pri) e0ti i ibtP ikit`i t¶t e eni SAKY neaq Avte` b bvgÄy Kiv nbj /</i></p> <p><i>USFDA KZR Abtgi` Z bq Ges BNF-G AŠ-fy bq neaq Avte` b bvgÄy Kiv hbZ cti /</i></p>	<p><i>Abtgi` b Kiv nbj /</i></p> <p><i>cōqirb bvB neaq Avte` b bvgÄy Kiv nbj /</i></p> <p><i>cōqirb bvB neaq Avte` b bvgÄy Kiv nbj /</i></p> <p><i>cōqirb bvB neaq Avte` b bvgÄy Kiv nbj /</i></p> <p><i>4(Pri) e0ti i ibtP ikit`i t¶t e eni SAKY neaq Avte` b bvgÄy Kiv nbj /</i></p> <p><i>USFDA KZR Abtgi` Z bq Ges BNF-G AŠ-fy bq neaq Avte` b bvgÄy Kiv nbj /</i></p>	

<i>bs</i>	<i>cōZKvi tKi bvg</i>	<i>Jl̄tai bvg I tR̄tbwK bvg</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigij 56 Zg mfvi imxvši</i>	<i>mfvi imxvš-</i>
Square Pharmaceuticals Ltd., Pabna	m)	<b>Alvimopan 12 mg Capsule</b>  Alvimopan INN 12 mg	Indicated to accelerate the time to gastrointestinal recovery following partial large or small bowel resection in patients at least 18 years of age. It is a µ-opioid receptor antagonist that prevents the peripheral GI effects of opioids on motility and secretion without reversing the central analgesia provided.	New	USFDA	<i>cōqirb bvB neaq Avte`b bvgÄy Kiv hwbZ c̄ti  </i>	<i>cōqirb bvB neaq Avte`b bvgÄy Kiv nbj  </i>
	n)	<b>Simethicone 80 mg Chewable Tablet</b>  Simethicone 100 DC grade Ph. Gr. (60%) 133.333 mg (eq. to Simethicone USP 80 mg)	Flatulence, abdominal distension, fullness, gas and windy colics. It is used for relief of the painful symptoms of excess gas in the digestive tract. Such gas is frequently caused by excessive swallowing of air or by eating foods that disagree.	67 mg/ml Paediatric Drops		<i>USFDA KZR.Abjgw` Z bq Ges BNF-G Aš-fg` bq neaq Avte`b bvgÄy Kiv hwbZ c̄ti  </i>	<i>USFDA KZR.Abjgw` Z bq Ges BNF-G Aš-fg` bq neaq Avte`b bvgÄy Kiv nbj  </i>
	o)	<b>Calcium 500 mg + Vitamin D<sub>3</sub> 440 IU/Sachet</b>  Calcium Carbonate BP 1250 mg (eq. to 500 mg Elemental Calcium) + Colecalciferol BP (Dry Vitamin D <sub>3</sub> ) 4.40 mg (eq. to 440 IU Vitamin D <sub>3</sub> )	Correction of Vitamin D and calcium combined deficiency in elderly people. Vitamin D and calcium supplementation as an adjunct to specific therapy for osteoporosis treatment in patients with established, or at high risk of Vitamin D and calcium combined deficiencies.	Calcium 500 mg + Vit. D <sub>3</sub> 200IU Tablet.	BNF-57	<i>cōqirb bvB neaq Avte`b bvgÄy Kiv hwbZ c̄ti  </i>	<i>Abjgy` b Kiv nbj  </i>

<i>bs</i>	<i>cōZKvi tKi bvg</i>	<i>Jl̄tai bvg I tR̄tbwK bvg</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-KigwJi 56 Zg mfvi imxvšl</i>	<i>mfvi imxvš-</i>
Square Pharmaceuticals Ltd., Pabna	p) <b>Pizotifen 0.25mg/5 ml Syrup</b>  Pizotifen Maleate BP 7.30 mg (eq. to Pizotifen 5 mg) /100 ml	Anorexia of somatic or psychogenic origin in underweight patients, supplementary to the treatment of underlying condition such as infectious or parasitic diseases, chronic diarrhea, anorexia nervosa or depressive states in the elderly.	500 mcg & 1.5 mg Tablet		USFDA KZR.Abtgwi Z bq Ges BNF-G Aš-fy bq neavq Avte`b bvgAý Kiv hvBtZ cüti	USFDA KZR.Abtgwi Z bq Ges BNF-G Aš-fy bq neavq Avte`b bvgAý Kiv nBj	
	q) <b>Flucytosine 500 mg Capsule</b>  Flucytosine BP 500 mg	Treatment of serious infections caused by susceptible strains of Candida and/or Cryptococcus, Candida: Septicemia, endocarditis and urinary system infections have been effectively treated with flucytosine.	New	USFDA	Abtgi`b Kiv hvBtZ cüti	Abtgi`b Kiv nBj	
	r) <b>Glimepiride 6 mg Tablet</b>  Glimepiride USP 6 mg	It is indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus.	1 mg, 2 mg & 4 mg Tablet	USFDA	cōqirb bvB neavq Avte`b bvgAý Kiv hvBtZ cüti	cōqirb bvB neavq Avte`b bvgAý Kiv nBj	
	s) <b>Glimepiride 8 mg Tablet</b>  Glimepiride USP 8 mg	-do-	-do-	USFDA	cōqirb bvB neavq Avte`b bvgAý Kiv hvBtZ cüti	cōqirb bvB neavq Avte`b bvgAý Kiv nBj	
	t) <b>Carvedilol Phosphate 10 mg CR Capsule</b>  Carvedilol Phosphate Controlled Release Pellets 20% w/w Ph. Gr. 50 mg contains : Carvedilol Phosphate INN 10 mg	Hypertension, heart failure and myocardial infarction	3.125 mg, 6.25 mg, 12.50 mg & 25 mg Tablet	USFDA	cōqirb bvB neavq Avte`b bvgAý Kiv hvBtZ cüti	cōqirb bvB neavq Avte`b bvgAý Kiv nBj	

<i>bs</i>	<i>cōZKvi tKi bvg</i>	<i>Jl̄tai bvg I tR̄tbwK bvg</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avtē` bKvix cōE</i> BNF or USFDA Reference	<i>tUKibK'ij mve-KigwJi 56 Zg mfvi m̄xv̄S̄</i>	<i>mfvi m̄xv̄S̄-</i>
	Square Pharmaceuticals Ltd., Pabna	<p>u) <b>Carvedilol Phosphate 20 mg CR Capsule</b> Carvedilol Phosphate Controlled Release Pellets 20% w/w Ph. Gr. 100 mg Contains : Carvedilol Phosphate INN 20 mg</p> <p>v) <b>Carvedilol Phosphate 40 mg CR Capsule</b> Carvedilol Phosphate Controlled Release Pellets 20% w/w Ph. Gr. 200 mg Contains : Carvedilol Phosphate INN 40 mg</p> <p>w) <b>Carvedilol Phosphate 80 mg CR Capsule</b> Carvedilol Phosphate Controlled Release Pellets 20% w/w Ph. Gr. 400 mg Contains : Carvedilol Phosphate INN 80 mg</p> <p>x) <b>Gabapentin 250 mg/5ml Syrup</b> Gabapentin USP 5 gm/100 ml</p> <p>y) <b>Amlodipine 5 mg + Hydrochlorothiazide 12.5 mg + Valsartan 160 mg Tablet</b> Amlodipine Besilate BP 7.00 mg eq. to 5 mg Amlodipine + Hydrochlorothiazide BP 12.50 mg + Valsartan USP 160 mg</p> <p>z) <b>Amlodipine 5 mg + Hydrochlorothiazide 25 mg + Valsartan 160 mg Tablet</b> Amlodipine Besilate BP 7.00 mg eq. to 5 mg Amlodipine + Hydrochlorothiazide BP 25 mg + Valsartan USP 160 mg</p>	Hypertension, heart failure and myocardial infarction  -do-  -do-  Postherpetic Neuralgia and Epilepsy.  Treatment of hypertension.  Treatment of hypertension.	3.125 mg, 6.25 mg, 12.50 mg & 25 mg Tablet  -  -  300 mg & 600 mg Tab.	USFDA  USFDA  USFDA  USFDA  New	<i>cōqiqRb bvb weavq Avte`b bvgAij Kiv hbBtZ cvti  </i>  <i>cōqiqRb bvb weavq Avte`b bvgAij Kiv hbBtZ cvti  </i>  <i>cōqiqRb bvb weavq Avte`b bvgAij Kiv hbBtZ cvti  </i>  <i>Abfgi`b Kiv hbBtZ cvti  </i>  <i>GB ḡT̄v cōqiqRb bvb weavq m̄m̄m̄-Gi 238 Zg mfviq Avte`b bvgAij Kii evi m̄xv̄S̄-enj iLV hbBtZ cvti  </i>	<i>cōqiqRb bvb weavq Avte`b bvgAij Kiv nbj  </i>  <i>cōqiqRb bvb weavq Avte`b bvgAij Kiv nbj  </i>  <i>cōqiqRb bvb weavq Avte`b bvgAij Kiv nbj  </i>  <i>Abfgi`b Kiv nbj  </i>  <i>GB ḡT̄v cōqiqRb bvb weavq Avte`b bvgAij Kiv nbj  </i>

<i>bs</i>	<i>cōZKvi tKi bvg</i>	<i>Jl̄tai bvg I tR̄tbwK bvg</i>	<i>ib̄t` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt̄e` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK̄yj mve-KiugwJi 56 Zg mfvi m̄xvš-</i>	<i>mfvi m̄xvš-</i>
Square Pharmaceuticals Ltd., Pabna	aa)	<b>Amlodipine 10 mg + Hydrochlorothiazide 12.50 mg + Valsartan 160 mg Tablet</b>  Amlodipine Besilate BP 14.00 mg eq. to 10 mg Amlodipine + Hydrochlorothiazide BP 12.50 mg + Valsartan USP 160 mg	Treatment of hypertension.	New	USFDA	<i>GB gr̄iv cōqyRb bvB neaq m̄mim-Gi 238 Zg mfviq Avt̄e` b bvgAiy Kiiv evi m̄xvš- enij iVLv hvBtZ cv̄ti  </i>	<i>GB gr̄iv cōqyRb bvB neaq Avt̄e` b bvgAiy Kiiv nBj  </i>
	ab)	<b>Amlodipine 10 mg + Hydrochlorothiazide 25 mg + Valsartan 160 mg Tablet</b>  Amlodipine Besilate BP 14.00 mg eq. to 10 mg Amlodipine + Hydrochlorothiazide BP 25 mg + Valsartan USP 160 mg	Treatment of hypertension.	New	USFDA	<i>GB gr̄iv cōqyRb bvB neaq m̄mim-Gi 238 Zg mfviq Avt̄e` b bvgAiy Kiiv evi m̄xvš- enij iVLv hvBtZ cv̄ti  </i>	<i>GB gr̄iv cōqyRb bvB neaq Avt̄e` b bvgAiy Kiiv nBj  </i>
	ac)	<b>Amlodipine 10 mg + Hydrochlorothiazide 25 mg + Valsartan 320 mg Tablet</b>  Amlodipine Besilate BP 14.00 mg eq. to 10 mg Amlodipine + Hydrochlorothiazide BP 25 mg + Valsartan USP 320 mg	Treatment of hypertension.	New	USFDA	<i>GB gr̄iv cōqyRb bvB neaq m̄mim-Gi 238 Zg mfviq Avt̄e` b bvgAiy Kiiv evi m̄xvš- enij iVLv hvBtZ cv̄ti  </i>	<i>GB gr̄iv cōqyRb bvB neaq Avt̄e` b bvgAiy Kiiv nBj  </i>
	ad)	<b>Amlodipine BP 5mg + Olmesartan Medoxomil 40mg Tablet</b>  Amlodipine Besilate BP 7.00 mg eq. to Amlodipine BP 5mg + Olmesartan Medoxomil INN 40mg	Treatment of hypertension.	New	USFDA	<i>Ab̄gr̄` b Kiiv hvBtZ cv̄ti  </i>	<i>Ab̄gr̄` b Kiiv nBj  </i>
	ae)	<b>Amlodipine BP 10mg + Olmesartan Medoxomil 20mg Tablet</b>  Amlodipine Besilate BP 14.00 mg eq. to Amlodipine BP 10mg + Olmesartan Medoxomil INN 20mg	Treatment of hypertension.	New	USFDA	<i>GB gr̄iv cōqyRb bvB neaq m̄mim-Gi 238 Zg mfviq Avt̄e` b bvgAiy Kiiv evi m̄xvš- enij iVLv hvBtZ cv̄ti  </i>	<i>GB gr̄iv cōqyRb bvB neaq Avt̄e` b bvgAiy Kiiv nBj  </i>

<i>bs</i>	<i>cōZKvi tKi bvg</i>	<i>Jl̄tai bvg I tR̄tbw̄K bvg</i>	<i>ib̄t` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt̄e` bKvix cōĒ BNF or USFDA Reference</i>	<i>tUKibK̄yj mve-Kiugw̄Ji 56 Zg mfvi m̄xv̄šl</i>	<i>mfvi m̄xv̄š-</i>
Square Pharmaceuticals Ltd., Pabna	af) <b>Amlodipine BP 10mg + Olmesartan Medoxomil 40mg Tablet</b>  Amlodipine Besilate BP 14.00 mg eq. to Amlodipine BP 10mg + Olmesartan Medoxomil INN 40mg	Treatment of hypertension.	New	USFDA	<i>GB gr̄iv cōqirRb bvB weavq m̄mim-Gi 238 Zg mfvq Avt̄e` b bvgĀy Kiv evi m̄xv̄š- enj i vLv h̄BtZ cīti  </i>	<i>GB gr̄iv cōqirRb bvB weavq Avt̄e` b bvgĀy Kiv nBj  </i>	
	ag) <b>Ascorbic Acid 1000mg + Elemental Zinc 10 mg Effervescent Tablet</b>  Ascorbic Acid BP 1000 mg + Zinc Sulphate Monohydrate USP 27.45 mg (eq. to 10 mg Elemental Zinc)	To maintain body immune system.	New		<i>GB gr̄iv cōqirRb bvB weavq m̄mim-Gi 238 Zg mfvq Avt̄e` b bvgĀy Kiv evi m̄xv̄š- enj i vLv h̄BtZ cīti  </i>	<i>GB gr̄iv cōqirRb bvB weavq Avt̄e` b bvgĀy Kiv nBj  </i>	
	ah) <b>Atomoxetine 10 mg Tablet</b>  Atomoxetine Hydrochloride INN 11.40 mg (eq. to Atomoxetine 10 mg)	Attention deficit hyperactivity disorder (initiated by a specialist physician experienced in managing the condition).	New	BNF-57 Page 217	<i>Abtgr̄` b Kiv h̄BtZ cīti  </i>	<i>Abtgr̄` b Kiv nBj  </i>	
	ai) <b>Zuclopentixol Acetate 50 mg/ml Injection Ampoule</b>  Zuclopentixol Acetate BP 50 mg/ml	Short term management of acute psychosis, mania, or exacerbations of chronic psychosis.	New	BNF-57 Page-197	<i>cōqirRb bvB weavq Avt̄e` b bvgĀy Kiv h̄BtZ cīti  </i>	<i>cōqirRb bvB weavq Avt̄e` b bvgĀy Kiv nBj  </i>	
	aj) <b>Zuclopentixol Decanoate 200 mg/ml Injection Ampoule</b>  Zuclopentixol Acetate BP 200 mg/ml	-do-	New	BNF-57	<i>Abtgr̄` b Kiv h̄BtZ cīti  </i>	<i>Abtgr̄` b Kiv nBj  </i>	

<i>bs</i>	<i>cōZKvi tKi bug</i>	<i>Jl̄tai bug I tR̄tbuiK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Aute` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigij 56 Zg mfvi imxviši</i>	<i>mfvi imxviš-</i>
	Square Pharmaceuticals Ltd., Pabna	<p>ak) <b>Light Liquid Paraffin 6 gm + White Soft Paraffin 15 gm/100 gm Cream</b>  Light Liquid Paraffin BP 6 gm + White Soft Paraffin BP 15 gm/100 gm</p> <p>al) <b>Tapentadol 50 mg Tablet</b>  Tapentadol Hydrochloride INN 58.245 mg eq. to 50 mg Tapentadol</p> <p>am) <b>Tapentadol 75 mg Tablet</b>  Tapentadol Hydrochloride INN 87.367 mg eq. to 75 mg Tapentadol</p> <p>an) <b>Tapentadol 100 mg Tablet</b>  Tapentadol Hydrochloride INN 116.49 mg eq. to 100 mg Tapentadol</p> <p>ao) <b>Doripenem 500 mg / Vial IV Injection</b>  Doripenem Monohydrate INN 521.403mg (eq. to Doripenem 500mg) /Vial</p> <p>ap) <b>Nitrofurantoin 25 mg/5ml Oral Suspension</b>  Nitrofurantoin BP 25 mg/5ml</p>	<p>To treat dry skin conditions.</p> <p>Indicated for the relief of moderate to severe acute pain in patients 18 years of age or older.</p> <p>-do-</p> <p>-do-</p> <p>It is indicated in the treatment of complicated Intra-abdominal Infections, Complicated UTI, including pyelonephritis and Nosocomial pneumonia and early-onset VAP(Ventilator-associated pneumonia).</p> <p>Urinary tract Infection due to susceptible strains of escherichiacoli, enterococci, staphylococcus aureus.</p>	<p>New</p> <p>New</p> <p>New</p> <p>New</p> <p>New</p>	<p>USFDA</p> <p>USFDA</p> <p>USFDA</p> <p>USFDA</p> <p>USFDA BNF-58</p> <p>USFDA BNF-58</p>	<p><i>Abfgi` b Kiv h̄BtZ cīti  </i></p>	<p><i>Abfgi` b Kiv nBj  </i></p>

<i>bs</i>	<i>cōZKvi tKi bvg</i>	<i>Jl̄tai bvg I tR̄tbwK bvg</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avtē` bKvix cōĒ BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigij 56 Zg mfvi m̄xvši</i>	<i>mfvi m̄xvš-</i>
	Square Pharmaceuticals Ltd., Pabna	<p>aq) <b>Piperacillin 3 gm + Tazobactam 375 mg/Vial IV Injection</b> Piperacillin Sodium USP 3.135 gm (eq. to Piperacillin 3 gm) + Tazobactam Sodium INN 0.404 gm (eq. to Tazobactam 375 mg)/Vial</p> <p>ar) <b>Piperacillin 2 gm + Tazobactam 250 mg/Vial IV Injection</b> Piperacillin Sodium USP 2.09 gm (eq. to Piperacillin 2 gm) + Tazobactam Sodium INN 0.269 gm (eq. to Tazobactam 250 mg)/Vial</p> <p>as) <b>Amlodipine 5 mg + Atorvastatin 20 mg FC Tablet</b> Amlodipine Besilate BP 7 mg eq. to Amlodipine 5 mg + Atorvastatin Calcium INN 20.72 mg eq. to Atorvastatin 20 mg</p> <p>at) <b>Ambrisentan 5 mg Tablet</b> Ambrisentan INN 5 mg</p> <p>au) <b>Ambrisentan 10 mg Tablet</b> Ambrisentan INN 10 mg</p>	<p>Lower respiratory tract infections, Urinary tract infections, intra abdominal infections, uncomplicated skin and skin structure infections</p> <p>-do-</p> <p>Hyperlipidemia with hypertension or angina.</p> <p>It is indicated for the treatment of patients with pulmonary arterial hypertension (PAH) classified as WHO functional class II and III, to improve exercise. Efficacy has been shown in idiopathic PAH (IPAH) and PAH associated with connective tissue disease.</p> <p>-do-</p>	<p>New</p> <p>New</p> <p>5 mg + 10 mg Tablet</p> <p>New</p> <p>New</p>	<p>USFDA</p> <p>USFDA BNF-58 Page-301</p> <p>USFDA</p> <p>BNF-58 Page : 96 USFDA</p> <p>BNF-58 Page : 96 USFDA</p>	<p>cōqiRb bvB neaq Avtē` b bvgĀy Kiv h̄BtZ c̄ti  </p> <p>Abtgv` b Kiv h̄BtZ c̄ti  </p> <p>GB ḡTv cōqiRb bvB neaq m̄m̄m-Gi 238 Zg mfviq Avtē` b bvgĀy Kiv eri m̄xvš-enyj ivLv h̄BtZ c̄ti  </p> <p>cōqiRb bvB neaq Avtē` b bvgĀy Kiv h̄BtZ c̄ti  </p> <p>cōqiRb bvB neaq Avtē` b bvgĀy Kiv h̄BtZ c̄ti  </p>	<p>cōqiRb bvB neaq Avtē` b bvgĀy Kiv nbj  </p> <p>Abtgv` b Kiv nbj  </p> <p>GB ḡTv cōqiRb bvB neaq Avtē` b bvgĀy Kiv nbj  </p> <p>cōqiRb bvB neaq Avtē` b bvgĀy Kiv nbj  </p> <p>cōqiRb bvB neaq Avtē` b bvgĀy Kiv nbj  </p>

<i>bs</i>	<i>cōZKvi tKi bvg</i>	<i>Jl̄tai bvg I tR̄tbw̄K bvg</i>	<i>ib̄t` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt̄e` bKvix cōE</i> BNF or USFDA Reference	<i>tUKibK̄yj mve-Kigw̄Uj 56 Zg mfvi m̄x̄iš̄</i>	<i>mfvi m̄x̄iš̄-</i>
	Square Pharmaceuticals Ltd., Pabna	av) <b>Guanethidine Monosulfate 10 mg Tablet</b>  Guanethidine Monosulphate BP 10 mg	It is indicated for the treatment of moderate and severe hypertension, either alone or as an adjunct, and for the treatment of renal hypertension, including that secondary to pyelonephritis, renal amyloidosis, and renal artery stenosis	New	USFDA	<i>cōq̄Rb b̄B m̄araq Avt̄e` b bvḡĀy Kiv h̄B̄tZ c̄t̄i  </i>	<i>cōq̄Rb b̄B m̄araq Avt̄e` b bvḡĀy Kiv nBj  </i>
		aw) <b>Guanethidine Monosulfate 25 mg Tablet</b>  Guanethidine Monosulphate BP 25 mg	-do-	New	USFDA	<i>cōq̄Rb b̄B m̄araq Avt̄e` b bvḡĀy Kiv h̄B̄tZ c̄t̄i  </i>	<i>cōq̄Rb b̄B m̄araq Avt̄e` b bvḡĀy Kiv nBj  </i>
		ax) <b>Moxonidine 0.20 mg Tablet</b>  Moxonidine BP 0.20 mg	Mild to moderate essential hypertension	New	BNF-58 Page 99	<i>cōq̄Rb b̄B m̄araq Avt̄e` b bvḡĀy Kiv h̄B̄tZ c̄t̄i  </i>	<i>cōq̄Rb b̄B m̄araq Avt̄e` b bvḡĀy Kiv nBj  </i>
		ay) <b>Moxonidine 0.30 mg Tablet</b>  Moxonidine BP 0.30 mg	-do-	New	BNF-58 Page 99	<i>cōq̄Rb b̄B m̄araq Avt̄e` b bvḡĀy Kiv h̄B̄tZ c̄t̄i  </i>	<i>cōq̄Rb b̄B m̄araq Avt̄e` b bvḡĀy Kiv nBj  </i>
		az) <b>Moxonidine 0.40 mg Tablet</b>  Moxonidine BP 0.40 mg	-do-	New	BNF-58 Page 99	<i>cōq̄Rb b̄B m̄araq Avt̄e` b bvḡĀy Kiv h̄B̄tZ c̄t̄i  </i>	<i>cōq̄Rb b̄B m̄araq Avt̄e` b bvḡĀy Kiv nBj  </i>
05.	Square Pharma Ltd., Gazipur	a) <b>Anhydrous Caffeine 30 mg/3 ml Injection</b>  Anhydrous Caffeine BP 30 mg/3 ml	Indicated for the short term treatment of apnea of prematurity.	New		USFDA KZR.Abtgw̄ Z bq Ges BNF-G Aš-f̄y bq m̄araq Avt̄e` b bvḡĀy Kiv h̄B̄tZ c̄t̄i	USFDA KZR.Abtgw̄ Z bq Ges BNF-G Aš-f̄y bq m̄araq Avt̄e` b bvḡĀy Kiv nBj

<i>bs</i>	<i>cōZKvičKi bug</i>	<i>Jl̄ai bug I tRtbičK bug</i>	<i>ibt`Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt`bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigij 56 Zg mfvi imxviši</i>	<i>mfvi imxviš-</i>
06.	Opsonin Pharma Ltd.	a) <b>Ondansetron 16 mg Suppository</b>  Ondansetron Hydrochloride USP 19.955 mg (eq. to Ondansetron 16 mg)	Indicated for the management of nausea and vomiting induced by cytotoxic chemotherapy and radiotherapy for the prevention of post operative nausea and vomiting in adults and for diarrhea induced nausea and vomiting.	4 mg & 8 mg Tablet 2mg/ml Injection	BNF-58 Page : 230	<i>Abfgi`b Kiv hvBtZ cti /</i>	<i>Abfgi`b Kiv nBj /</i>
		b) <b>Pizotifen 5 mg/100 ml Elixir</b>  Pizotifen Maleate BP 7.269 mg (eq. to Pizotifen 5 mg)	Indicated for the prophylactic treatment of vascular headaches of the migraine type such as classical migraine, common migraine and cluster headache.	500 mcg & 1.5 mg Tablet	BNF-57 (Page 248)	<i>cōqirb bvB neaq Avte`b bvgAý Kiv hvBtZ cti /</i>	<i>cōqirb bvB neaq Avte`b bigAý Kiv nBj /</i>
07.	Pharmasia Limited	a) <b>Multivitamin-Multimineral Sprinkle</b>  Each sachet contain one gram micronutrient mix including Vitamin A (Vitamin A Acetate) USP 0.400 mg, Vitamin E USP 5.000 mg Vitamin D USP 5.000 mcg Vitamin C ( Ascorbic Acid) USP 30.000 mg Vitamin B12 (Cyanocobalamin) BP 0.900 mcg, Vitamin B1 (Thiamine Mononitrate) BP 0.500 mg, Vitamin B2 (Riboflavin) BP 0.500 mg, Vitamin B6 (Pyridoxine Hydrochloride) BP 0.500 mg, Folic Acid BP 0.150 mg, Nicotinic Acid BP 6.000 mg, Iron (as encapsulated Ferrous Fumarate) BP 12.500 mg, Zinc ( as Gluconate) USP 4.100 mg, Copper USP 0.560 mg Iodine (as Potassium Iodide) USP 90.000 mcg Selenium USP 17.000 mcg	It is indicated to meet the increased demands of vitamins and minerals for growing children. This preparation is also indicated for the prevention and treatment of vitamins and minerals deficiencies and lack of appetite of children.	New		<i>AvciZZt -MZ ivLv hvBtZ cti /</i>	<i>AvciZZt -MZ ivLv nBj /</i>

<i>bs</i>	<i>cōZKvi tKi bug</i>	<i>Jl̄tai bug I tR̄tbuiK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Aute` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kiguij 56 Zg mfvi imxviši</i>	<i>mfvi imxviš-</i>
	Pharmasia Limited	b) <b>Ramelteon 8 mg Tablet</b>  Ramelteon INN 8 mg	Indicated for the treatment of insomnia characterized by difficulty with sleep onset.	New	USFDA approved (July 22, 2005)	<i>Abtgi`b Kiv hvBtZ c̄ti /</i>	<i>Abtgi`b Kiv nBj /</i>
		c) <b>Lithium 120 mg Tablet</b>  Lithium Orotate INN 120 mg	Mood stabilizer, treatment of bipolar disorder since, unlike most other mood altering drugs, they counteract both mania and depression.	New		<i>USFDA KZR.Abtgiw` Z bq Ges BNF-G Aš-fq bq neavq Aute`b bigAiy Kiv hvBtZ c̄ti /</i>	<i>Psychiatric Patients- t` i PiKrmiq e`envi nq neavq Jlaiui Safety, Efficacy m=utKgZigZ c̄vtbi Rb Aa'icK tMjvg i e'vbx, c̄wi Pij K, RvZiq gibimK -r- Bb=ilJDU Ges Aa'vcK I qmRDj Avj g tPšaj, RvZiq gibimK -r- Bb=ilJDU Gi ibKU tc̄Y Kiv hvBtZ c̄ti /</i>
		d) <b>Atorvastatin 10 mg + Niacin 375 mg Capsule</b>  Atorvastatin Calcium trihydrate INN 10.82 mg equ. to Atorvastatin 10 mg + Niacin USP 375 mg	Antihyperlipidemia & Prevention of Cardiovascular disease.	New		<i>USFDA KZR.Abtgiw` Z bq Ges BNF-G Aš-fq bq neavq Aute`b bigAiy Kiv hvBtZ c̄ti /</i>	<i>USFDA KZR.Abtgiw` Z bq Ges BNF-G Aš-fq bq neavq Aute`b bigAiy Kiv nBj /</i>
		e) <b>Atorvastatin 10 mg + Niacin 500 Capsule</b>  Atorvastatin Calcium trihydrate INN 10.82 mg equ. to Atorvastatin 10 mg + Niacin USP 500 mg Capsule	Antihyperlipidemia & Prevention of Cardiovascular disease.	New		<i>USFDA KZR.Abtgiw` Z bq Ges BNF-G Aš-fq bq neavq Aute`b bigAiy Kiv hvBtZ c̄ti /</i>	<i>USFDA KZR.Abtgiw` Z bq Ges BNF-G Aš-fq bq neavq Aute`b bigAiy Kiv nBj /</i>
		f) <b>Metaxalone 400 mg + Diclofenac Potassium 50 mgTablet</b>  Metaxalone INN 400 mg + Diclofenac Potassium BP 50 mg Tablet	Primary dysmenorrhea, mild to moderate pain, sign & symptoms of osteoarthritis & rheumatoid arthritis.	New		<i>USFDA KZR.Abtgiw` Z bq Ges BNF-G Aš-fq bq neavq Aute`b bigAiy Kiv hvBtZ c̄ti /</i>	<i>USFDA KZR.Abtgiw` Z bq Ges BNF-G Aš-fq bq neavq Aute`b bigAiy Kiv nBj /</i>
		g) <b>Paracetamol 500 mg + Pamabrom 25 mg Tablet</b>  Paracetamol BP 500 mg + Pamabrom USP25 mg Tablet	Premenstrual & Menstrual pain.	New		<i>USFDA KZR.Abtgiw` Z bq Ges BNF-G Aš-fq bq neavq Aute`b bigAiy Kiv hvBtZ c̄ti /</i>	<i>USFDA KZR.Abtgiw` Z bq Ges BNF-G Aš-fq bq neavq Aute`b bigAiy Kiv nBj /</i>

<i>bs</i>	<i>cōZKvi tKi bug</i>	<i>Jl̄tai bug I tR̄tbuiK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigilji 56 Zg mfvi imxvišl</i>	<i>mfvi imxviš-</i>
	Pharmasia Limited	<p>h) <b>Doxylamine Succinate 10 mg + Pyridoxine HCl 10 mg Extended release Tablet</b>  Doxylamine Succinate USP 10 mg + Pyridoxine HCl BP 10 mg Tablet</p> <p>i) <b>Alvimopan 12 mg Capsule</b>  Alvimopan INN 12 mg</p> <p>j) <b>Mesalazine 250 mg ER Capsule</b>  Mesalazine Extended Release Granules 300 mg Ph Gr contains Mesalazine USP 250 mg</p> <p>k) <b>Mesalazine 500 mg ER Capsule</b>  Mesalazine Extended Release Granules 600 mg Ph Gr contains Mesalazine USP 500 mg</p> <p>l) <b>Halobetasol 0.05% Ointment</b>  Halobetasol Propionate (Microfine) INN 0.050 gm/100 gm</p> <p>m) <b>Ceftibuten 400 mg Capsule</b>  Ceftibuten Dihydrate INN 435.085 mg eq. to Ceftibuten 400 mg</p> <p>n) <b>Ceftibuten 90 mg/5 ml Powder for Suspension</b>  Ceftibuten Dihydrate INN 1.958 gm eq. to Ceftibuten 1.80 gm/100 ml</p>	<p>Nausea &amp; vomiting during pregnancy.</p> <p>Indicated in patients to avoid postoperative ileum following partial large or small bowel resection with primary anastomosis.</p> <p>Ulcerative colitis.</p> <p>Ulcerative colitis.</p> <p>For the treatment of variety of skin conditions (e.g., eczema, dermatitis, allergies, rash).</p> <p>Acute bacterial Exacerbations of Chronic Bronchitis, Acute bacterial Otitis Media, Pharyngitis, Pharynitis &amp; Pneumonia.</p> <p>-do-</p>	<p>New</p> <p>New</p> <p>New</p> <p>New</p> <p>New</p> <p>New</p>	<p>USFDA approved</p> <p>USFDA</p> <p>BNF-58 (page-54) &amp; USFDA</p> <p>BNF-58 (page-54) &amp; USFDA</p> <p>USFDA</p> <p>USFDA</p>	<p>GB Kibkb cōqirb bvB neaq Avte`b bigAij Kiv hbZ citi  </p> <p>cōqirb bvB neaq Avte`b bigAij Kiv hbZ citi  </p> <p>Abtgv`b Kiv hbZ citi  </p>	<p>GB Kibkb cōqirb bvB neaq Avte`b bigAij Kiv nBj  </p> <p>cōqirb bvB neaq Avte`b bigAij Kiv nBj  </p> <p>Abtgv`b Kiv nBj  </p>

<i>bs</i>	<i>cōZKvi tKi bug</i>	<i>Jl̄tai bug I tR̄tbuiK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Aute` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigilji 56 Zg mfvi imxviši</i>	<i>mfvi imxviš-</i>
	Pharmasia Limited	<p>o) <b>Amlodipine 5 mg + Telmisartan 80 mg Tablet</b> Amlodipine besilate BP 6.94 mg equ. to Amlodipine 5 mg &amp; Telmisartan INN 80 mg Tablet</p> <p>p) <b>Amlodipine 5 mg + Telmisartan 40 mg Tablet</b> Amlodipine besilate BP 6.94 mg equ. to Amlodipine 5 mg &amp; Telmisartan INN 40 mg</p> <p>q) <b>Chlorpheniramine Maleate 2 mg + Codeine 10 mg Extended release suspension</b> Chlorpheniramine maleate (as Polistirex complex) BP 2 mg &amp; Codeine (as Polistirex complex) USP 10 mg / 5 ml</p> <p>r) <b>Chlorpheniramine Maleate 4 mg + Codeine 20 mg/5 ml Extended release suspension</b> Chlorpheniramine maleate (as Polistirex complex) BP 4 mg &amp; Codeine (as Polistirex complex) USP 20 mg / 5 ml</p> <p>s) <b>Chlorpheniramine Maleate 4 mg + Codeine 10 mg Tablet</b> Chlorpheniramine maleateBP 4 mg &amp; Codeine USP 10 mg Tablet</p> <p>t) <b>Chlorpheniramine Maleate 4 mg + Codeine 20 mg Tablet</b> Chlorpheniramine maleate BP 4 mg &amp; Codeine USP 20 mg Tablet</p> <p>u) <b>Fluocinolone Acetonide 0.01% + Hydroquinolone 4% + Tretinoin 0.05% Cream</b> Fluocinolone acetonide (Microfine) BP 0.01%+ Hydroquinolone (Microfine) USP 4%+Tretinoin (Microfine) BP 0.05% Cream</p>	Treatment of hypertension, chronic stable angina & Vasospastic angina.	New	USFDA	<i>GB Kirtbkb c̄qiRb bvB neaq Aute` b bigAy Kiv hbZ c̄ti  </i>	<i>GB Kirtbkb c̄qiRb bvB neaq Aute` b bigAy Kiv nBj  </i>
			Treatment of hypertension, chronic stable angina & Vasospastic angina.	New	USFDA	<i>Abtgr` b Kiv hbZ c̄ti  </i>	<i>Abtgr` b Kiv nBj  </i>
			Cough, runny nose, sneezing, itching of the nose or throat, and itchy watery eyes due to hay fever.	New		<i>USFDA KZR Abtgr` Z bq Ges BNF-G AŠ-fy bq neaq Aute` b bigAy Kiv hbZ c̄ti  </i>	<i>USFDA KZR Abtgr` Z bq Ges BNF-G AŠ-fy bq neaq Aute` b bigAy Kiv nBj  </i>
			Cough, runny nose, sneezing, itching of the nose or throat, and itchy watery eyes due to hay fever.	New	USFDA	<i>c̄qiRb bvB neaq Aute` b bigAy Kiv hbZ c̄ti  </i>	<i>c̄qiRb bvB neaq Aute` b bigAy Kiv nBj  </i>
			Cough, runny nose, sneezing, itching of the nose or throat, and itchy watery eyes due to hay fever.	New		<i>c̄qiRb bvB neaq Aute` b bigAy Kiv hbZ c̄ti  </i>	<i>c̄qiRb bvB neaq Aute` b bigAy Kiv nBj  </i>
			Cough, runny nose, sneezing, itching of the nose or throat, and itchy watery eyes due to hay fever.	New	USFDA	<i>c̄qiRb bvB neaq Aute` b bigAy Kiv hbZ c̄ti  </i>	<i>c̄qiRb bvB neaq Aute` b bigAy Kiv nBj  </i>
			Moderate to severe melasma of the face.	New	USFDA	<i>Abtgr` b Kiv hbZ c̄ti  </i>	<i>Abtgr` b Kiv nBj  </i>

<i>bs</i>	<i>cōZKvičKi bug</i>	<i>Jl̄ai bug I tR̄bniK bug</i>	<i>ibt`Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Aute`bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-KiugUj 56 Zg mfvi imxvši</i>	<i>mfvi imxvš-</i>
08.	Monicopharma Ltd.	a) <b>Cefixime Granules 50 mg Sachet</b>  Cefixime Granules USP 50 mg/Sachet	Treatment of urinary tract infections, upper and lower respiratory tract infections, acute otitis media, gonococcal urethritis and enteric fever.	100 mg, 200 mg & 400 mg Capsule		USFDA KZR Abtgw Z bq Ges BNF-G Aš-fy bq neavq Aute`b bugAý Ki v hvBtZ cüti	USFDA KZR Abtgw Z bq Ges BNF-G Aš-fy bq neavq Aute`b bugAý Ki v nBj
		b) <b>Ezetimibe 10 mg + Simvastatin 20 mg Tablet</b>  Ezetimibe INN 10 mg + Simvastatin USP 20 mg	Indicated for the reduction of elevated total-C and LDL-C levels in adult and adolescent (10 to 70 years of age) patients with HoFH. Patients may also receive adjunctive treatments.	New	USFDA BNF-55	KirAbkb ActqiqRbxq neavq Wim-Gi 238 Zg mfviq Aute`b bugAý Ki v evi imxvš- enj i Lv hvBtZ cüti	KirAbkb ActqiqRbxq neavq Aute`b bugAý Ki v nBj
		c) <b>Febuxostat 40 mg Tablet</b>  Febuxostat INN 40 mg	Indicated for the long term management of hyperuricemia in patients with gout. It is not recommended for use in the treatment of asymptomatic hyperuricemia.	New	USFDA	Abtgv`b Ki v hvBtZ cüti	Abtgv`b Ki v nBj
		d) <b>Varenicline 0.5 mg Tablet</b>  Varenicline INN 0.5 mg	Indicated as an aid to smoking cessation.	New	USFDA BNF-56	Abtgv`b Ki v hvBtZ cüti	Abtgv`b Ki v nBj
09.	Popular Pharmaceuticals Ltd.	a) <b>Medroxyprogesterone Acetate 10 mg Tablet</b>  Medroxyprogesterone Acetate USP 10 mg	For the treatment of menstrual irregularities and other disorders caused by inadequate endocrine secretion of the corpus luteum hormone.	150mg/ml Injection & 5 mg Tablet	BNF-56 Page-398	Abtgv`b Ki v hvBtZ cüti	Abtgv`b Ki v nBj

<i>bs</i>	<i>cōZKvi tKi bug</i>	<i>Jl̄tai bug I tR̄tbuiK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Ait̄e` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigilji 56 Zg mfvi m̄xv̄ši</i>	<i>mfvi m̄xv̄š-</i>
Popular Pharmaceuticals Ltd.	b) <b>Pregabalin 25 mg Capsule</b>  Pregabalin INN 25 mg	Pregabalin is an antiepileptic used as an adjunct in the treatment of partial seizures with or without secondary generalization. It is also used in the treatment of generalized anxiety disorder and neuropathic pain.	50 mg, 75 mg, 100 mg & 150 mg Capsule	USFDA BNF-57 Page-253	<i>Abfgi` b Kiv h̄BtZ c̄ti i /</i>	<i>Abfgi` b Kiv nBj /</i>	
	c) <b>Oseltamivir 30 mg Capsule</b>  Oseltamivir Phosphate INN 39.42 mg (eq. to 30 mg Oseltamivir)	Oseltamivir is indicated for the treatment of uncomplicated acute illness due to influenza infection in patients 1 year and older who have been symptomatic for no more than 2 days. Oseltamivir is indicated for the prophylaxis of influenza in patients 1 year and older.	75 mg, 45 mg Capsule 60 mg/5 ml Powder for Suspension	USFDA BNF-57 Page-350	<i>Abfgi` b Kiv h̄BtZ c̄ti i /</i>	<i>Abfgi` b Kiv nBj /</i>	
	d) <b>Oral Rehydration Salts (ORS) in Sachet</b>  Sodium Chloride BP 0.65 gm + Potassium Chloride BP 0.375 gm + Sodium Citrate, Dihydrate (Trisodium Citrate, Dihydrate) BP 0.725 gm + Anhydrous Glucose (Glucose, Anhydrous) BP 3.375 gm/Sachet	Used to treat and prevent dehydration (the loss of too much water) due to diarrhoea and gastroenteritis in children and adults.	Sod. Chloride 1.30 g Pot. Chloride 0.75 g Sod. Citrate 1.45 g Anhydrous Glucose 6.75 g		<i>c̄qiqRb b̄B m̄earq Ait̄e` b b̄gĀy Kiv h̄BtZ c̄ti i /</i>	<i>c̄qiqRb b̄B m̄earq Ait̄e` b b̄gĀy Kiv nBj /</i>	
	e) <b>Ebastine 10mg Fast Dissolving Tablet</b>  Ebastine BP 10mg	For the symptomatic treatment of Seasonal and perennial allergic rhinitis, Idiopathic chronic urticaria.	10 mg Tablet		<i>c̄qiqRb b̄B m̄earq m̄imim-Gi 238 Zg mfviq Ait̄e` b b̄gĀy Kiv evi m̄xv̄š- enij ivLv h̄BtZ c̄ti i /</i>	<i>c̄qiqRb b̄B m̄earq Ait̄e` b b̄gĀy Kiv nBj /</i>	
	f) <b>Ebastine 5mg/5 ml Syrup</b>  Ebastine BP 0.10 gm/100 ml	-do-	10 mg Tablet		<i>USFDA KZR Abfgi` Z bq Ges BNF-G AS-fy bq m̄earq Ait̄e` b b̄gĀy Kiv h̄BtZ c̄ti i /</i>	<i>USFDA KZR Abfgi` Z bq Ges BNF-G AS-fy bq m̄earq Ait̄e` b b̄gĀy Kiv nBj /</i>	

<i>bs</i>	<i>cōZKvičKi bvg</i>	<i>Jl̄tai bvg I tR̄tbvčK bvg</i>	<i>ibt`Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt`bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigilji 56 Zg mfvi imxviši</i>	<i>mfvi imxviš-</i>
10.	UniMed & UniHealth Manufacturers Ltd.	<p>a) <b>Calcium 500 + Vitamin D 400 IU Tablet</b>  Calcium Carbonate USP 1250 mg (eq. to 500 mg Elemental Calcium) + Cholecalciferol USP 4 mg (eq. to Vitamin D 400 IU)</p> <p>b) <b>Calcium 600 mg + Vitamin D 400 IU + Mineral Supplement Tablet</b>  Calcium Carbonate USP 1500 mg (eq. to 600 mg Elemental Calcium) + Cholecalciferol USP 4 mg (Vitamin D 400 IU) + Magnesium Oxide USP 66.34 mg (Magnesium 40 mg) + Zinc Oxide BP 9.34 mg (Zinc 7.5 mg) + Cupric Oxide Ph. Gr 1.25 mg (Copper 1 mg) + Manganese Sulfate USP 4.94 mg (Manganese 1.8 mg) + Boron Citrate Ph. Gr. 4 mg (Boron 0.25 mg)</p> <p>c) <b>Prazosin 5 mg Extended Release Tablet</b>  Prazosin Hydrochloride BP 5.476 mg (eq. to 5 mg Prazosin)</p> <p>d) <b>Prazosin 2.5 mg Extended Release Tablet</b>  Prazosin Hydrochloride BP 2.738 mg (eq. to 2.5 mg Prazosin)</p> <p>e) <b>Dipyridamole 200 mg + Aspirin 25 mg Extended Release capsule</b>  Dipyridamole USP 200 mg + Aspirin USP 25 mg</p>	<p>Prevention of Vitamin D and calcium deficiency</p> <p>Treatment to Vitamin D, calcium and mineral deficiency.</p> <p>For the treatment of all grades of essential hypertension and of all grades of secondary hypertension of varied aetiology.</p> <p>-do-</p> <p>Indicated to reduce the risk of stroke in patients who have had transient ischemia of the brain or completed ischemic stroke due to thrombosis.</p>	<p>500 mg + 200IU Tab.</p> <p></p> <p>1 mg &amp; 2 mg Tab.</p> <p>1 mg &amp; 2 mg Tab.</p> <p>New</p>	<p>BNF-56 Page : 534</p> <p></p> <p></p> <p></p> <p>BNF-56 Page : 132 USFDA</p>	<p>cōqirb bvB neaq Avte`b bvgAý Kiv hbZ cüti  </p> <p>USFDA KZR Abtgw Z bq Ges BNF-G Aš-fy bq neaq Avte`b bvgAý Kiv hbZ cüti  </p> <p>USFDA KZR Abtgw Z bq Ges BNF-G Aš-fy bq neaq Avte`b bvgAý Kiv hbZ cüti  </p> <p>USFDA KZR Abtgw Z bq Ges BNF-G Aš-fy bq neaq Avte`b bvgAý Kiv hbZ cüti  </p> <p>cōqirb bvB neaq Avte`b bvgAý Kiv hbZ cüti  </p>	<p>Abfgi`b Kiv nBj  </p> <p>USFDA KZR Abtgw Z bq Ges BNF-G Aš-fy bq neaq Avte`b bvgAý Kiv nBj  </p> <p>USFDA KZR Abtgw Z bq Ges BNF-G Aš-fy bq neaq Avte`b bvgAý Kiv nBj  </p> <p>USFDA KZR Abtgw Z bq Ges BNF-G Aš-fy bq neaq Avte`b bvgAý Kiv nBj  </p> <p>cōqirb bvB neaq Avte`b bvgAý Kiv nBj  </p>

<i>bs</i>	<i>cōZKvi tKi bug</i>	<i>Jl̄tai bug I tR̄tbw̄K bug</i>	<i>ib̄t` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt̄e` bKvix cōE</i> BNF or USFDA Reference	<i>tUKibK̄yj mve-KigilJi 56 Zg mfvi imx̄išl</i>	<i>mfvi imx̄iš-</i>
	UniMed & UniHealth Manufacturers Ltd.	f) <b>Sibutramine Hydrochloride Monohydrate 15 mg Capsule</b>  Sibutramine Hydrochloride Monohydrate INN 15 mg	Management of obesity	5 mg Tab. & 5 mg Cap.	BNF-57 Page : 221 USFDA	<i>cōqiqRb bvB neaq Avte`b bigAý Kiv hvBtZ cv̄ti  </i>	<i>cōqiqRb bvB neaq Avte`b bigAý Kiv nBj  </i>
		g) <b>Sibutramine Hydrochloride Monohydrate 10 mg Capsule</b>  Sibutramine Hydrochloride Monohydrate INN 10 mg	Management of obesity	5 mg Tab. & 5 mg Cap.	BNF-57 Page : 221 USFDA	<i>cōqiqRb bvB neaq Avte`b bigAý Kiv hvBtZ cv̄ti  </i>	<i>cōqiqRb bvB neaq Avte`b bigAý Kiv nBj  </i>
		h) <b>Tadalafil 10mg Tablet</b>  Tadalafil INN 10mg	It is indicated for erectile dysfunction.	New	BNF 58 USFDA	<i>Acēenvi nBtZ cv̄ti neaq mvgiRK tc̄lvcU netePbv Kii qv Avte`b -MZ ivLv hvBtZ cv̄ti  </i>	<i>Acēenvi nBtZ cv̄ti neaq mvgiRK tc̄lvcU netePbv Kii qv Avte`b -MZ ivLv nBj  </i>
		i) <b>Tadalafil 20mg Tablet</b>  Tadalafil INN 20mg	-do-	New	BNF 58 USFDA	<i>Acēenvi nBtZ cv̄ti neaq mvgiRK tc̄lvcU netePbv Kii qv Avte`b -MZ ivLv hvBtZ cv̄ti  </i>	<i>Acēenvi nBtZ cv̄ti neaq mvgiRK tc̄lvcU netePbv Kii qv Avte`b -MZ ivLv nBj  </i>
		j) <b>Sildenafil 25mg Tablet</b>  Sildenafil Citrate INN 35.121 mg (eq. to Sildenafil 25 mg)	It is indicated for pulmonary arterial hypertension, erectile dysfunction.	New	BNF 58 USFDA	<i>Acēenvi nBtZ cv̄ti neaq mvgiRK tc̄lvcU netePbv Kii qv Avte`b -MZ ivLv hvBtZ cv̄ti  </i>	<i>Acēenvi nBtZ cv̄ti neaq mvgiRK tc̄lvcU netePbv Kii qv Avte`b -MZ ivLv nBj  </i>
		k) <b>Sildenafil 50mg Tablet</b>  Sildenafil Citrate INN 70.241 mg (eq. to Sildenafil 50 mg)	-do-	New	BNF 58 USFDA	<i>Acēenvi nBtZ cv̄ti neaq mvgiRK tc̄lvcU netePbv Kii qv Avte`b -MZ ivLv hvBtZ cv̄ti  </i>	<i>Acēenvi nBtZ cv̄ti neaq mvgiRK tc̄lvcU netePbv Kii qv Avte`b -MZ ivLv nBj  </i>
		l) <b>Sildenafil 100mg Tablet</b>  Sildenafil Citrate INN 140.482 mg (eq. to Sildenafil 100 mg)	-do-	New	BNF 58 USFDA	<i>Acēenvi nBtZ cv̄ti neaq mvgiRK tc̄lvcU netePbv Kii qv Avte`b -MZ ivLv hvBtZ cv̄ti  </i>	<i>Acēenvi nBtZ cv̄ti neaq mvgiRK tc̄lvcU netePbv Kii qv Avte`b -MZ ivLv nBj  </i>

<i>bs</i>	<i>cōZKvičKi bug</i>	<i>Jl̄tai bug I tR̄tbniK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Aute` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigilji 56 Zg mfvi imxvišl</i>	<i>mfvi imxviš-</i>
11.	Beacon Pharmaceuticals Ltd.	a) <b>Paricalcitol 1 mcg Capsule</b>  Paricalcitol USP 1 mcg	Treatment of secondary hyperparathyroidism associated with chronic renal insufficiency.	New	BNF-57 Page 543	<i>Abfgv` b Kiv hvBtZ ciči /</i>	<i>Abfgv` b Kiv nBj /</i>
	b) <b>Paricalcitol 2 mcg Capsule</b>  Paricalcitol USP 2 mg	-do-	New	BNF-57 Page 543	<i>Abfgv` b Kiv hvBtZ ciči /</i>	<i>Abfgv` b Kiv nBj /</i>	
	c) <b>Vinorelbine 10 mg/ ml Injection Vial</b>  Vinorelbine Tartrate USP 13.85 mg (eq. to 10 mg Vinorelbine)/ 1 ml	Indicated as a single agent or in combination with cisplatin for the first line treatment of ambulatory patients with unresectable, advance non-small cell lung cancer.	New	BNF-57 (Page 474) USFDA	<i>Abfgv` b Kiv hvBtZ ciči /</i>	<i>Abfgv` b Kiv nBj /</i>	
	d) <b>Vinorelbine 50 mg/ 5 ml Injection Vial</b>  Vinorelbine Tartrate USP 69.27 mg (eq. to 50 mg Vinorelbine)/ 5 ml	-do-	New		<i>Abfgv` b Kiv hvBtZ ciči /</i>	<i>Abfgv` b Kiv nBj /</i>	
	e) <b>Tinzaparin Sodium 3500 IU/0.35 ml Injection Prefilled Syringe</b>  Tinzaparin Sodium INN 3500 IU/0.35 ml	For the prevention of thromboembolic events including deep vein thrombosis, in patients undergoing general and orthopaedic surgery, treatment of deep vein thrombosis, treatment of pulmonary embolism and prevention of coagulation of blood in extra corporeal circulation, such as haemodialysis.	New	BNF 57 Page : 126	<i>Abfgv` b Kiv hvBtZ ciči /</i>	<i>Abfgv` b Kiv nBj /</i>	
	f) <b>Palonosetron 0.25 mg/5 ml Injection, Vial</b>  Palonosetron Hydrochloride INN 0.28 mg (eq. to 0.25 mg of Palonosetron)/5 ml	Moderately emetogenic cancer chemotherapy, Highly emetogenic cancer chemotherapy, Prevention of postoperative nausea and vomiting for up to 24 hours following surgery.	New	USFDA	<i>Abfgv` b Kiv hvBtZ ciči /</i>	<i>Abfgv` b Kiv nBj /</i>	

<i>bs</i>	<i>cōZKvičKi bug</i>	<i>Jl̄ai bug I tRtbičK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigilji 56 Zg mfvi imxvišl</i>	<i>mfvi imxviš-</i>
	Beacon Pharmaceuticals Ltd.	<p>g) <b>Palonsetron 0.075 mg /1.5 ml Injection, Vial</b>  Palonosetron Hydrochloride INN 0.084 mg (eq. to 0.075 mg of Palonosetron)/1.5 ml</p> <p>h) <b>Oprelvekin 5 mg/ml Lyophilized Injection, Vial</b>  Oprelvekin INN 5 mg/ml</p> <p>i) <b>Prasugrel 5 mg Tablet</b>  Prasugrel Hydrochloride INN 5.49 mg ( Eq.to Prasugrel 5 mg )</p> <p>j) <b>Prasugrel 10 mg Tablet</b>  Prasugrel Hydrochloride INN 10.98 mg ( Eq.to Prasugrel 10 mg )</p> <p>k) <b>Paclitaxel 300 mg/50 ml Injection</b>  Paclitaxel USP 300 mg/50 ml vial</p>	<p>Moderately emetogenic cancer chemotherapy, Highly emetogenic cancer chemotherapy, Prevention of postoperative nausea and vomiting for up to 24 hours following surgery.</p> <p>Indicated for the prevention of severe thrombocytopenia and the reduction of the need for platelet transfusion following myelosuppressive chemotherapy in adult patients with nonmyeloid malignancies who are at high risk of severe thrombocytopenia.</p> <p>It is indicated to reduce the rate of thrombotic cardiovascular events in patients with acute coronary syndrome, patient with unstable angina or non-ST-elevation myocardial infarction.</p>	<p>New</p> <p>New</p> <p>New</p> <p>New</p>	<p>-do-</p> <p>USFDA</p> <p>USFDA</p> <p>USFDA</p> <p>30 mg/Vial &amp; 100 mg Vial BNF 57 (Page 84)</p>	<p><i>Abfgv` b Kiv hvBtZ cüti  </i></p>	<p><i>Abfgv` b Kiv nBj  </i></p>

<i>bs</i>	<i>cōZKvičKi bug</i>	<i>Jl̄tač bug I tRtbičK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Ače` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigilji 56 Zg mfvi wmxvišl</i>	<i>mfvi wmxviš-</i>
	Beacon Pharmaceuticals Ltd.	<p>l) <b>Ifosfamide 1 gm/Vial Lyophilized Powder for Injection</b>  Ifosfamide USP 1 gm/Vial</p> <p>m) <b>Ifosfamide 2 gm/Vial Lyophilized Powder for Injection</b>  Ifosfamide USP 2 gm/Vial</p> <p>n) <b>Trastuzumab 440 mg/Vial Lyophilized Powder for Injection</b>  Trastuzumab INN 440 mg/Vial</p> <p>o) <b>Fluconazole 50 mg/25 ml Infusion</b>  Fluconazole USP 50 mg/25 ml</p> <p>p) <b>Fluconazole 200 mg/100 ml Infusion</b>  Fluconazole USP 200 mg/100 ml</p> <p>q) <b>Fluconazole 400 mg/200 ml Infusion</b>  Fluconazole USP 400 mg/200 ml</p>	<p>It is indicated for third line chemotherapy of germ cell testicular cancer. It should ordinarily be used in combination with a prophylactic agent for hemorrhagic cystitis.</p> <p>-do-</p> <p>It is indicated for adjuvant treatment of HER2 Over expressing node positive or node negative breast cancer</p> <p>It is indicated for the treatment of Oropharyngeal and esophageal candidiasis. In open noncomparative studies of relatively small numbers of patients, fluconazole also effective for the treatment of candida urinary tract infections, peritonitis, and systemic candida infections including candidemia, disseminated candidiasis, and pneumonia</p> <p>-do-</p> <p>-do-</p>	<p>New</p> <p>New</p> <p>New</p> <p>50 mg, 150 mg &amp; 200 mg Cap. 3mg/ml Eye Drops 50mg./5 ml Powder for Suspension</p> <p>-do-</p> <p>-do-</p>	<p>BNF-57 Page : 465</p> <p>BNF-57 Page : 465</p> <p>USFDA BNF-57 Page : 485</p> <p>USFDA (2mg/ml) BNF : 57 Page : 330</p> <p>USFDA</p> <p>USFDA</p>	<p><i>Abfgi` b Kiv hBtZ cti i /</i></p>	<p><i>Abfgi` b Kiv nBj /</i></p>

<i>bs</i>	<i>cōZKvičKi bug</i>	<i>Jl̄ai bug I tR̄bniK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigilji 56 Zg mfvi m̄xvišl</i>	<i>mfvi m̄xviš-</i>
Beacon Pharmaceuticals Ltd.	r) <b>Pemetrexed 100 mg/ ml Lyophilized Powder for Injection</b>  Pemetrexed Disodium USP 108 mg (eq. to 100 mg Pemetrexed)/1 ml Vial	Anti-Cancer.	New	USFDA	<i>Abfgi`b Kiv h̄BtZ c̄ti  </i>	<i>Abfgi`b Kiv nBj  </i>	
		Anti-Cancer.	New	USFDA	<i>Abfgi`b Kiv h̄BtZ c̄ti  </i>	<i>Abfgi`b Kiv nBj  </i>	
	l) <b>Eptifibatide 75 mg/100 ml Injection</b>  Eptifibatide INN 75 mg/100 ml Vial	It is used usually in combination with aspirin and heparin, in the management of unstable angina and in patients undergoing coronary angioplasty and stenting procedures.	New	BNF-57 Page : 134 USFDA	<i>Abfgi`b Kiv h̄BtZ c̄ti  </i>	<i>Abfgi`b Kiv nBj  </i>	
12. Aristopharma Ltd.	a) <b>Lornoxicam 4 mg Tablet</b>  Lornoxicam INN 4 mg	Lornoxicam is an oxicam derivative is a NSAID. It is used in musculoskeletal and joint disorders such as osteoarthritis and rheumatoid arthritis.	New		USFDA KZR <i>Abfgi` Z bq Ges BNF-G Aš-fy bq meaq Avte`b bigÄy Kiv h̄BtZ c̄ti  </i>	USFDA KZR <i>Abfgi` Z bq Ges BNF-G Aš-fy bq meaq Avte`b bigÄy Kiv nBj  </i>	
	b) <b>Lornoxicam 8 mg Tablet</b>  Lornoxicam INN 8 mg	-do-	New		USFDA KZR <i>Abfgi` Z bq Ges BNF-G Aš-fy bq meaq Avte`b bigÄy Kiv h̄BtZ c̄ti  </i>	USFDA KZR <i>Abfgi` Z bq Ges BNF-G Aš-fy bq meaq Avte`b bigÄy Kiv nBj  </i>	
	c) <b>Paracetamol 500 mg + Anhydrous Codeine Phosphate 8 mg Tablet</b>  Paracetamol BP 500 mg + Codeine Phosphate BP 8.18 mg (eq. to Anhydrous Codeine Phosphate 8 mg)	Indicated for the relief of mild to moderate pain such as; headache, period pains, neuralgia, toothache, rheumatic pains and symptoms of cold and influenza.	New	BNF-58	<i>c̄qyRb bvB meaq Avte`b bigÄy Kiv h̄BtZ c̄ti  </i>	<i>c̄qyRb bvB meaq Avte`b bigÄy Kiv nBj  </i>	

<i>bs</i>	<i>cōZKvičKi bug</i>	<i>Jl̄ai bug I tR̄bniK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigilji 56 Zg mfvi imxviši</i>	<i>mfvi imxviš-</i>
	Aristopharma Ltd.	d) <b>Paracetamol 500 mg + Anhydrous Codeine Phosphate 15 mg Tablet</b>  Paracetamol BP 500 mg + Codeine Phosphate BP 15.345 mg (eq. to Anhydrous Codeine Phosphate 15 mg)	Indicated for the relief of mild to moderate pain such as; headache, period pains, neuralgia, toothache, rheumatic pains and symptoms of cold and influenza.	New	BNF-58	cōqirb bvB neaq Avte`b bigAjy Kiv hbZ cti /	cōqirb bvB neaq Avte`b bigAjy Kiv nbj /
		e) <b>Paracetamol 500 mg + Anhydrous Codeine Phosphate 30 mg Tablet</b>  Paracetamol BP 500 mg + Codeine Phosphate BP 30.690 mg (eq. to Anhydrous Codeine Phosphate 30 mg)	-do-	New	BNF-58	cōqirb bvB neaq Avte`b bigAjy Kiv hbZ cti /	cōqirb bvB neaq Avte`b bigAjy Kiv nbj /
		f) <b>Doxofylline 400 mg Tablet</b>  Doxofylline INN 400 mg	Indicated for the treatment of bronchial asthma and chronic obstructive pulmonary disease (COPD) in adults.	New		USFDA KZR.Abtgwi Z bq Ges BNF-G Aš-fj bq neaq Avte`b bigAjy Kiv hbZ cti /	USFDA KZR.Abtgwi Z bq Ges BNF-G Aš-fj bq neaq Avte`b bigAjy Kiv nbj /
		g) <b>Aceclofenac 200 mg Sustained release Tablet</b>  Aceclofenac BP 200 mg	Indicated for the treatment of pain and inflammation in rheumatoid arthritis, osteoarthritis and ankylosing spondylitis.	100 mg Tablet		USFDA KZR.Abtgwi Z bq Ges BNF-G Aš-fj bq neaq Avte`b bigAjy Kiv hbZ cti /	USFDA KZR.Abtgwi Z bq Ges BNF-G Aš-fj bq neaq Avte`b bigAjy Kiv nbj /
		h) <b>Sodium Chloride 50mg/ml Eye Drops</b>  Sodium Chloride BP 5 gm/100 ml	Indicated for the temporary relief of corneal edema.	0.9 gm/100 ml Eye Drops		USFDA KZR.Abtgwi Z bq Ges BNF-G Aš-fj bq neaq Avte`b bigAjy Kiv hbZ cti /	USFDA KZR.Abtgwi Z bq Ges BNF-G Aš-fj bq neaq Avte`b bigAjy Kiv nbj /
		i) <b>Sodium Chloride 50 mg/gm Eye Ointment</b>  Sodium Chloride BP 5 gm/100 gm	-do-	-do-		USFDA KZR.Abtgwi Z bq Ges BNF-G Aš-fj bq neaq Avte`b bigAjy Kiv hbZ cti /	USFDA KZR.Abtgwi Z bq Ges BNF-G Aš-fj bq neaq Avte`b bigAjy Kiv nbj /

<i>bs</i>	<i>cōZKvi tKi bvg</i>	<i>Jl̄tai bvg I tR̄tbwK bvg</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-KigwJi 56 Zg mfvi m̄xvšl</i>	<i>mfvi m̄xvš-</i>
	Aristopharma Ltd	j) <b>Trifluridine 10 mg/ml Eye Drops</b>  Trifluridine USP 1 gm/100 ml	It is used to treat eye infections caused by the herpes simplex virus, which can lead to swelling or ulcers in the eyelids or cornea.	New	USFDA	<i>Abfgi` b Kiv h̄BtZ c̄ti  </i>	<i>Abfgi` b Kiv nBj  </i>
		k) <b>Paracetamol 500 mg + Diphenhydramine HCl 25 mg Tablet</b>  Paracetamol BP 500 mg + Diphenhydramine HCl USP 25 mg	For the relief of the symptoms associated with colds and influenza, mild to moderate pain such as headaches, migraine, rheumatic and muscle pain.	New		<i>Kir̄bkb Ac̄qirRbxq neaq Wimim-Gi 238 Zg mfvq Avt` b bvgÄy Kiv evi m̄xvš- enjy i vLv h̄BtZ c̄ti  </i>	<i>Kir̄bkb Ac̄qirRbxq neaq Avt` b bvgÄy Kiv nBj  </i>
		l) <b>Paracetamol 120 mg + Diphenhydramine Hydrochloride 12.50 mg/5 ml Syrup</b>  Paracetamol BP 2.40 gm + Diphenhydramine Hydrochloride BP 0.250 gm/100 ml	-do-	New		<i>USFDA KZR.Abstgw` Z bq Ges BNF-G AS-fy` bq neaq Avte` b bvgÄy Kiv h̄BtZ c̄ti  </i>	<i>USFDA KZR.Abstgw` Z bq Ges BNF-G AS-fy` bq neaq Avt` b bvgÄy Kiv nBj  </i>
		m) <b>Bisoprolol Hemifumarate 2.5 mg + Hydrochlorothiazide 6.25 mg Tablet</b>  Bisoprolol Hemifumarate USP 2.5 mg + Hydrochlorothiazide BP 6.25 mg	Indicated for the treatment of hypertension.	New	USFDA	<i>c̄qyRb bvB neaq Avte` b bvgÄy Kiv h̄BtZ c̄ti  </i>	<i>c̄qyRb bvB neaq Avte` b bvgÄy Kiv nBj  </i>
		n) <b>Bisoprolol Hemifumarate 5 mg + Hydrochlorothiazide 6.25 mg Tablet</b>  Bisoprolol Hemifumarate USP 5 mg + Hydrochlorothiazide BP 6.25 mg	Indicated for the treatment of hypertension.	New	USFDA	<i>c̄qyRb bvB neaq Avte` b bvgÄy Kiv h̄BtZ c̄ti  </i>	<i>c̄qyRb bvB neaq Avte` b bvgÄy Kiv nBj  </i>
		o) <b>Bisoprolol Hemifumarate 10 mg + Hydrochlorothiazide 6.25 mg Tablet</b>  Bisoprolol Hemifumarate USP 10 mg + Hydrochlorothiazide BP 6.25 mg	Indicated for the treatment of hypertension.	New	USFDA	<i>c̄qyRb bvB neaq Avte` b bvgÄy Kiv h̄BtZ c̄ti  </i>	<i>c̄qyRb bvB neaq Avte` b bvgÄy Kiv nBj  </i>

<i>bs</i>	<i>cōZKvi tKi bvg</i>	<i>Jl̄tai bvg I tR̄tbwK bvg</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Aute` bKvix cōE</i> BNF or USFDA Reference	<i>tUKibK'ij mve-KigwJi 56 Zg mfvi imxvši</i>	<i>mfvi imxvš-</i>
Aristopharma Ltd.	p) Thiamine Nitrate 300 mg + Pyridoxine HCl 300 mg + Cyanocobalamin 1 mg + Vitamin E 100 IU Tablet  Thiamine Nitrate BP 300 mg + Pyridoxine Hydrochloride BP 300 mg + Cyanocobalamin BP 1 mg + Vitamin E USP 100 IU	Indicated where a deficiency of the relevant vitamins.	New		USFDA KZR.Abtgw` Z bq Ges BNF-G Aš-fy bq neavq Aute` b bvgAý Kiv hBtZ cüti	USFDA KZR.Abtgw` Z bq Ges BNF-G Aš-fy bq neavq Aute` b bvgAý Kiv nBj	
	q) <b>Bumetanide 5 mg Tablet</b>  Bumetanide USP 5 mg	Oedema associated with congestive heart failure, hepatic & renal disease including the nephrotic syndrome.	1 mg Tablet	BNF-58 Page-77	Abtgw` b Kiv hBtZ cüti	Abtgw` b Kiv nBj	
	r) <b>Bumetanide 2 mg/4 ml Injection Ampoule</b>  Bumetanide (Injectable grade) USP 2 mg/4ml	Oedema associated with congestive heart failure, hepatic & renal disease including the nephrotic syndrome.	1 mg Tablet	BNF-58 Page-77	Abtgw` b Kiv hBtZ cüti	Abtgw` b Kiv nBj	
	s) <b>Gliclazide 40 mg Tablet</b>  Gliclazide BP 40 mg	It is indicated for control of blood glucose in patients with non-insulin dependent diabetes mellitus whose hyperglycemia can not be controlled by diet alone.	30 mg & 80 mg Tablet		USFDA KZR.Abtgw` Z bq Ges BNF-G Aš-fy bq neavq Aute` b bvgAý Kiv hBtZ cüti	USFDA KZR.Abtgw` Z bq Ges BNF-G Aš-fy bq neavq Aute` b bvgAý Kiv nBj	
	t) <b>Rabeprazole Sodium 20 mg/Vial Lyophilized Injection (IV)</b>  Rabeprazole Sodium INN 20 mg/Vial	Rabeprazole IV is an alternative in patients for whom oral administration of rabeprazole is not indicated. It is indicated in the treatment of Active duodenal ulcer with bleeding or severe erosions, Active gastric ulcer with bleeding or severe erosions and Short-term treatment of erosive or ulcerative gastroesophageal reflux disease.	10 mg & 20 mg Tablet		USFDA KZR.Abtgw` Z bq Ges BNF-G Aš-fy bq neavq Aute` b bvgAý Kiv hBtZ cüti	USFDA KZR.Abtgw` Z bq Ges BNF-G Aš-fy bq neavq Aute` b bvgAý Kiv nBj	

<i>bs</i>	<i>cōZKvičKi bug</i>	<i>Jl̄ai bug I tRtbičK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Aute` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigilji 56 Zg mfvi imxvišl</i>	<i>mfvi imxviš-</i>
13.	Incepta Pharmaceuticals Ltd.,	a) <b>Prasugrel 5mg Tablet</b>  Prasugrel HCl INN 5.49mg eq. to Prasugrel 5mg	Indicated for the prevention of atherothrombotic events in patients with acute coronary syndrome	New	USFDA	<i>Abfgv` b Kiv hBtZ cti i /</i>	<i>Abfgv` b Kiv nBj /</i>
	b) <b>Prasugrel 10mg Tablet</b>  Prasugrel HCl INN 10.98mg eq. to Prasugrel 10mg	-do-	New	USFDA	<i>Abfgv` b Kiv hBtZ cti i /</i>	<i>Abfgv` b Kiv nBj /</i>	
	c) <b>Padimate O 8%+Avobenzone 2%+Oxybenzone 3%+Titanium Dioxide 2% Lotion</b>  Padimate O USP 8gm+Avobenzone USP 2gm+Oxybenzone USP 3gm+Titanium Dioxide BP 2gm/100ml	Sunblock to protect the skin from UVA and UVB rays, for people whose skin is abnormally sensitive to sunlight, for example due to radiotherapy or taking certain medicines eg. amiodarone, tetracycline antibiotics, or due to skin conditions such as solar urticaria, acute solar dermatitis, acute lupus erythematosus, herpes simplex, polymorphic light eruption, cutaneous albinism or vitiligo.	New	BNF-57, Page 644	<i>Abfgv` b Kiv hBtZ cti i /</i>	<i>Abfgv` b Kiv nBj /</i>	
	d) <b>Nitrofurantoin 25mg/5ml Suspension</b>  Nitrofurantoin BP 0.5gm/100ml	Indicated for the prophylaxis and treatment of infections of the genito-urinary tract due to susceptible bacteria.	100mg tablet	USFDA	<i>Abfgv` b Kiv hBtZ cti i /</i>	<i>Abfgv` b Kiv nBj /</i>	
	e) <b>Besifloxacin 0.6% Suspension of Eye Drops</b>  Besifloxacin HCl INN 0.663gm eq. to Besifloxacin 0.6gm/100ml	Indicated for the treatment of bacterial conjunctivitis.	New	USFDA	<i>Abfgv` b Kiv hBtZ cti i /</i>	<i>Abfgv` b Kiv nBj /</i>	

<i>bs</i>	<i>cōZKvičKi bug</i>	<i>Jl̄ai bug I tRtbičK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Aiče` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-KigilUi 56 Zg mfvi imxvišl</i>	<i>mfvi imxviš-</i>
	Incepta Pharmaceuticals Ltd.,	f) <b>Amphotericin B (Liposomal) 50mg/Vial Injection</b>  Amphotericin B (Liposomal) (as Sterile Lyophilised Powder) USP 50mg/Vial	It is indicated for the treatment of visceral leishmaniasis. Indicated for prophylaxis in liver transplant patients at risk of systemic Candida, Aspergillus and Cryptococcus infections, and for the treatment of systemic fungal infections caused by organisms susceptible to Amphotericin B. Amphotericin is indicated for empirical treatment of presumed fungal infections in febrile neutropaenic patients whose fever has failed to respond to broad spectrum antibiotic treatment..	New	BNF-58, Page 334	<i>Abfgi` b Kiv hvBtZ ciči /</i>	<i>Abfgi` b Kiv nBj /</i>
		g) <b>Conjugated Estrogens 1.25 mg Tablet</b>  Conjugated Estrogens USP 1.250 mg	Prevention of postmenopausal osteoporosis. Treatment of moderate to severe vasomotor symptoms associated with the menopause. Treatment of moderate to severe symptoms of vulvar and vaginal atrophy associated with the menopause.	Conjugated Estrogens 0.625 mg Tablet	USFDA	<i>Abfgi` b Kiv hvBtZ ciči /</i>	<i>c`ilUi Safety, Efficacy m=útkčgiq gj-vqtbj Rb- tUKibK'ij mve-KigilUi mfviq Dc-vctbj Rb- ejv nBj /</i>
		h) <b>Conjugated Estrogen 25 mg/vial Injection</b>  Conjugated Estrogen USP 25 mg/Vial (Sterile Lyophilized Powder)	-do-	-do-	USFDA	<i>Abfgi` b Kiv hvBtZ ciči /</i>	<i>Abfgi` b Kiv nBj /</i>

<i>bs</i>	<i>cōZKvi tKi bvg</i>	<i>Jl̄tai bvg I tR̄tbwK bvg</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-KiugUj 56 Zg mfvi m×vš-</i>	<i>mfvi m×vš-</i>
14.	Alco Pharma Ltd.	a) Ezetimibe 10 mg + Simvastatin 10 mg Tablet  Ezetimibe INN 10 mg + Simvastatin USP 10 mg	Primary Hypercholesterolemia or mixed hyperlipidemia.	New	USFDA	Kifbkb AcqRbq neaq wWmm-Gi 238 Zg mfvi Avte`b bvgAij Kii evi m×vš-enj ivLv hbTZ cti	Kifbkb AcqRbq neaq Avte`b bvgAij Kiv nbj
	b) Ezetimibe 10 mg + Simvastatin 20 mg Tablet  Ezetimibe INN 10 mg + Simvastatin USP 20 mg	-do-		New	USFDA BNF	Kifbkb AcqRbq neaq wWmm-Gi 238 Zg mfvi Avte`b bvgAij Kii evi m×vš-enj ivLv hbTZ cti	Kifbkb AcqRbq neaq Avte`b bvgAij Kiv nbj
	c) Loperamide 2mg + Simethicone 125mg Chewable Tablet  Loperamide HCl BP 2.153 mg eq. to Loperamide 2 mg + Simethicone USP 125mg	Treat Diarrhoea and Symptoms of gas.		New	USFDA	cqRb bvB neaq wWmm-Gi 238 Zg mfvi Avte`b bvgAij Kii evi m×vš-enj ivLv hbTZ cti	cqRb bvB neaq Avte`b bvgAij Kiv nbj
	d) Loperamide 2mg + Simethicone 125mg Tablet  Loperamide HCl BP 2.153 mg eq. to Loperamide 2 mg+ Simethicone USP 125mg	Treat Diarrhoea and Symptoms of gas.		New	USFDA BNF	cqRb bvB neaq wWmm-Gi 238 Zg mfvi Avte`b bvgAij Kii evi m×vš-enj ivLv hbTZ cti	cqRb bvB neaq Avte`b bvgAij Kiv nbj
	e) Acetaminophen 325 mg + Diphenhydramine HCl 12.50 mg + Phenylephrine HCl 5 mg Tablet  Acetaminophen USP 325 mg + Diphenhydramine Hydrochloride USP 12.50 mg + Phenylephrine Hydrochloride BP 5 mg	Temporarily relieves cold, flu symptom, minor aches and pain.		New		Kifbkb AcqRbq neaq wWmm-Gi 238 Zg mfvi Avte`b bvgAij Kii evi m×vš-enj ivLv hbTZ cti	Kifbkb AcqRbq neaq Avte`b bvgAij Kiv nbj
	f) Repaglinide 1mg+Metformin HCl 500mg Tablet  Repaglinide BP 1mg+Metformin HCl BP 500mg	Adjunct to diet and exercise to improve glycemic control in adults with type diabetes mellitus.		New	USFDA	cqRb bvB neaq wWmm-Gi 238 Zg mfvi Avte`b bvgAij Kii evi m×vš-enj ivLv hbTZ cti	cqRb bvB neaq Avte`b bvgAij Kiv nbj

<i>bs</i>	<i>cōZKvičKi bvg</i>	<i>Jlčai bvg I tRtbičK bvg</i>	<i>ibt`Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt`bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigilji 56 Zg mfvi m×vši</i>	<i>mfvi m×vš-</i>
	Alco Pharma Ltd.	g) <b>Repaglinide 2mg+Metformin HCl 500mg Tablet</b>  Repaglinide BP 2mg+Metformin HCl BP 500mg	Adjunct to diet and exercise to improve glycemic control in adults with type2 diabetes mellitus.	New	USFDA	<i>cōqiqRb bvB neavq m×vš-Gi 238 Zg mfviq Avte`b bvgAý Kiv evi m×vš- enyj ivLv hvBtZ cüti /</i>	<i>cōqiqRb bvB neavq Avte`b bvgAý Kiv nBj /</i>
15.	ACI Ltd.	a) <b>Dimercaprol 100 mg/ 2ml Injection Ampoule</b>  Dimercaprol BP 100mg/2 ml	It is indicated in the treatment of Arsenic,Gold & Mercury poisoning.	New	BNF-58	<i>cōqiqRb bvB neavq Avte`b bvgAý Kiv hvBtZ cüti /</i>	<i>cōqiqRb bvB neavq Avte`b bvgAý Kiv nBj /</i>
	b) <b>Dimercaprol 50 mg/ ml Injection Ampoule</b>  Dimercaprol BP 50mg/ml	Do	New			<i>Abtgv`b Kiv hvBtZ cüti /</i>	<i>Abtgv`b Kiv nBj /</i>
	c) <b>Doripenem 500 mg Injection Vial</b>  Doripenem Monohydrate INN 0.5214 gm (eq. to Doripenem 0.5 gm)/Vial	It is indicated in the treatment of Nosocomial Pneumonia, Complicated UTI, Complicated Intra-abdominal Infection.	New	USFDA BNF-58		<i>Abtgv`b Kiv hvBtZ cüti /</i>	<i>Abtgv`b Kiv nBj /</i>
	d) <b>Glycopyrrolate 0.2 mg/ml Injection Ampoule</b>  Glycopyrrolate USP 0.2mg/ml	Glycopyrrolate is used to reduce saliva, nasal,lung and stomach secretion and to help control heart rate during surgery.	New	USFDA		<i>Abtgv`b Kiv hvBtZ cüti /</i>	<i>Abtgv`b Kiv nBj /</i>
	e) <b>Prasugrel 5 mg Tablet</b>  Prasugrel Hydrochloride INN 5.489 mg (eq. to Prasugrel 5 mg)	Persugrel is indicated for prevention of Atherothrombotic events in patients with acute coronary syndrome eg unstable Angina, Miocardial Infarction.	New	BNF-58 USFDA		<i>Abtgv`b Kiv hvBtZ cüti /</i>	<i>Abtgv`b Kiv nBj /</i>
	f) <b>Prasugrel 10 mg Tablet</b>  Prasugrel Hydrochloride INN 10.978 mg (eq. to Prasugrel 10 mg)	-do-	New	BNF-58 USFDA		<i>Abtgv`b Kiv hvBtZ cüti /</i>	<i>Abtgv`b Kiv nBj /</i>

<i>bs</i>	<i>cōZKvičKi bug</i>	<i>Jlčai bug I tRčbičK bug</i>	<i>ibč Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avtč bKvix cōč BNF or USFDA Reference</i>	<i>tUKčbKčj mve-Kigilji 56 Zg mfvi mčxviši</i>	<i>mfvi mčxviš-</i>
ACI Ltd.	g) Isoproterenol Hydrochloride 0.2 mg/ml Injection Ampoule  Isoprenaline Hydrochloride USP 0.2mg/ml	It is indicated for 1. Mild or transient episodes of heart block that do not require electric shock or pacemaker therapy 2. Serious episodes of heart block 3.Bronchospasm occurring during anaesthesia.	New			IMZ ivLv hBtZ cti Ges cZōvbičK cčqRbčq tidiči Ymn cčvivq Avte`b Kiičevi Rb` ejv hBtZ cti /	IMZ ivLv nBj Ges cZōvbičK cčqRbčq tidiči Ymn cčvivq Avte`b Kiičevi Rb` ejv nBj /
	h) Rabeprazole Sodium 20 mg + Domperidone 30 mg Capsule  250 mg Blended pellets contains : Rabeprazole Sodium INN 20 mg + Domperidone BP 30 mg	It is indicated for 1. Dyspepsia 2. GERD 3. Nausia associated with peptic disorder 4. Post operative nausea & vomiting 5. Chronic gastritis.	New			USFDA KZR Abtgw Z bq Ges BNF-G Aš-fč bq neaq Avte`b bigAčy Kiv hBtZ cti /	USFDA KZR Abtgw Z bq Ges BNF-G Aš-fč bq neaq Avte`b bigAčy Kiv nBj /
	i) Esomeprazole Sodium 20 mg + Domperidone 30 mg Capsule  250 mg Blended pellets contains : Esomeprazole Sodium INN 20mg + Domperidone BP 30mg	It is indicated for 1. Dyspepsia 2. GERD 3. Nausia associated with peptic disorder 4. Post operative nausea & vomiting 5. Chronic gastritis.	New			USFDA KZR Abtgw Z bq Ges BNF-G Aš-fč bq neaq Avte`b bigAčy Kiv hBtZ cti /	USFDA KZR Abtgw Z bq Ges BNF-G Aš-fč bq neaq Avte`b bigAčy Kiv nBj /
	j) Dexamethasone 0.1% + Moxifloxacin 0.5% Eye Drops  Dexamethasone Sodium Phosphate BP 0.1316 Eq.to Dexamethasone 0.1 gm + Moxifloxacin HCl INN 0.545gm Eq. to Moxifloxacin 0.5 gm/100 ml	It is indicated for complication after surgery and external eye diseases, specially for the elderly.	New			IMZ ivLv hBtZ cti Ges cZōvbičK cčqRbčq tidiči Ymn cčvivq Avte`b Kiičevi Rb` ejv hBtZ cti /	IMZ ivLv nBj Ges cZōvbičK cčqRbčq tidiči Ymn cčvivq Avte`b Kiičevi Rb` ejv nBj /
	k) Doxapram HCl 100mg/5 ml IV Injection Ampoule  Doxapram HCl USP 100 mg/5 ml	Respiratory Stimulants & Pulmonary surfactants.	New	BNF 58 USFDA		Abtgw`b Kiv hBtZ cti /	Abtgw`b Kiv hBtZ cti /
	l) Sildenafil 20mg Tablet  Sildenafil Citrate INN 28.097 mg (eq. to Sildenafil 20 mg)	It is indicated for pulmonary arterial hypertension, erectile dysfunction.	New	BNF 58 USFDA		Ace`envi nBtZ cti neaq mgmRK tcčvU nečPbv Kiičvq Avte`b IMZ ivLv hBtZ cti /	Ace`envi nBtZ cti neaq mgmRK tcčvU nečPbv Kiičvq Avte`b IMZ ivLv nBj /

<i>bs</i>	<i>cōZKvičKi bug</i>	<i>Jl̄ai bug I tR̄bniK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigilj i 56 Zg mfvi imxviši</i>	<i>mfvi imxviš-</i>
ACI Ltd.	m) <b>Sildenafil 25mg Tablet</b>  Sildenafil Citrate INN 35.121 mg (eq. to Sildenafil 25 mg)	It is indicated for pulmonary arterial hypertension, erectile dysfunction.	New	BNF 58 USFDA	Ace`envi nBtZ c̄ti weaq mvgmRK tcljvcU wePbv Kui qv Avt` b -MZ ivLv h̄BtZ c̄ti /	Ace`envi nBtZ c̄ti weaq mvgmRK tcljvcU wePbv Kui qv Avt` b -MZ ivLv nbj /	
	n) <b>Sildenafil 50mg Tablet</b>  Sildenafil Citrate INN 70.241 mg (eq. to Sildenafil 50 mg)	-do-	New	BNF 58 USFDA	Ace`envi nBtZ c̄ti weaq mvgmRK tcljvcU wePbv Kui qv Avt` b -MZ ivLv h̄BtZ c̄ti /	Ace`envi nBtZ c̄ti weaq mvgmRK tcljvcU wePbv Kui qv Avt` b -MZ ivLv nbj /	
	o) <b>Sildenafil 100mg Tablet</b>  Sildenafil Citrate INN 140.482 mg (eq. to Sildenafil 100 mg)	-do-	New	BNF 58 USFDA	Ace`envi nBtZ c̄ti weaq mvgmRK tcljvcU wePbv Kui qv Avt` b -MZ ivLv h̄BtZ c̄ti /	Ace`envi nBtZ c̄ti weaq mvgmRK tcljvcU wePbv Kui qv Avt` b -MZ ivLv nbj /	
	p) <b>Tadalafil 2.5mg Tablet</b>  Tadalafil INN 2.5mg	It is indicated for erectile dysfunction.	New	BNF 58 USFDA	Ace`envi nBtZ c̄ti weaq mvgmRK tcljvcU wePbv Kui qv Avt` b -MZ ivLv h̄BtZ c̄ti /	Ace`envi nBtZ c̄ti weaq mvgmRK tcljvcU wePbv Kui qv Avt` b -MZ ivLv nbj /	
	q) <b>Tadalafil 5mg Tablet</b>  Tadalafil INN 5mg	-do-	New	BNF 58 USFDA	Ace`envi nBtZ c̄ti weaq mvgmRK tcljvcU wePbv Kui qv Avt` b -MZ ivLv h̄BtZ c̄ti /	Ace`envi nBtZ c̄ti weaq mvgmRK tcljvcU wePbv Kui qv Avt` b -MZ ivLv nbj /	
	r) <b>Tadalafil 10mg Tablet</b>  Tadalafil INN 10mg	-do-	New	BNF 58 USFDA	Ace`envi nBtZ c̄ti weaq mvgmRK tcljvcU wePbv Kui qv Avt` b -MZ ivLv h̄BtZ c̄ti /	Ace`envi nBtZ c̄ti weaq mvgmRK tcljvcU wePbv Kui qv Avt` b -MZ ivLv nbj /	
	s) <b>Tadalafil 20mg Tablet</b>  Tadalafil INN 20mg	-do-	New	BNF 58 USFDA	Ace`envi nBtZ c̄ti weaq mvgmRK tcljvcU wePbv Kui qv Avt` b -MZ ivLv h̄BtZ c̄ti /	Ace`envi nBtZ c̄ti weaq mvgmRK tcljvcU wePbv Kui qv Avt` b -MZ ivLv nbj /	
	t) <b>Cefaclor 250mg Chewable Tablet</b>  Cefaclor Monohydrate USP 262.245 mg (eq. to Cefaclor 250mg)	Infections due to Gram-positive and Gram-negative bacteria eg. Septicaemia, pneumonia, meningitis, peritonitis, U.T.I. etc.	250 mg & 500 mg Capsule & 375 mg Tablet	USFDA	c̄qirb bvB weaq Avt` b bigÄy Kiv h̄BtZ c̄ti /	c̄qirb bvB weaq Avt` b bigÄy Kiv nbj /	
	u) <b>Cefaclor 500mg Chewable Tablet</b>  Cefaclor Monohydrate USP 524.49 mg (eq. to Cefaclor 500 mg)	-do-			USFDA KZK Abfgw Z bq Ges BNF-G Aš-fy bq weaq Avt` b bigÄy Kiv h̄BtZ c̄ti /	USFDA KZK Abfgw Z bq Ges BNF-G Aš-fy bq weaq Avt` b bigÄy Kiv nbj /	

<i>bs</i>	<i>cōZKvičKi bug</i>	<i>Jl̄tai bug I tR̄tbuiK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kiugij 56 Zg mfvi m̄xvši</i>	<i>mfvi m̄xvš-</i>
	ACI Ltd.	v) <b>Amlodipine 10 mg + Valsartan 160 mg + Hydrochlorothiazide 12.50 mg Tablet</b>  Amlodipine Besilate BP 13.86 mg (eq. to Amlodipine 10 mg) + Valsartan USP 160 mg + Hydrochlorothiazide BP 12.5 mg	This combination is indicated for treatment of hypertension. (It is not indicated for initial therapy of hypertension).	New	USFDA	<i>GB ḡīv c̄l̄q̄Rb b̄B m̄avq m̄m̄m-Gi 238 Zg mfviq Avt` b bugĀy K̄i evi m̄xvš- enj iLv h̄B̄tZ c̄t̄i /</i>	<i>GB ḡīv c̄l̄q̄Rb b̄B m̄avq Avt` b bugĀy K̄i nBj /</i>
	w) <b>Amlodipine 10 mg + Valsartan 160 mg + Hydrochlorothiazide 25 mg Tablet</b>  Amlodipine Besilate BP 13.86 mg (eq. to Amlodipine 10 mg) + Valsartan USP 160 mg + Hydrochlorothiazide BP 25 mg	-do-	New	USFDA	<i>GB ḡīv c̄l̄q̄Rb b̄B m̄avq m̄m̄m-Gi 238 Zg mfviq Avt` b bugĀy K̄i evi m̄xvš- enj iLv h̄B̄tZ c̄t̄i /</i>	<i>GB ḡīv c̄l̄q̄Rb b̄B m̄avq Avt` b bugĀy K̄i nBj /</i>	
	x) <b>Amlodipine 10 mg + Valsartan 320 mg + Hydrochlorothiazide 25 mg Tablet</b>  Amlodipine Besilate BP 13.86 mg (eq. to Amlodipine 10 mg) + Valsartan USP 320 mg + Hydrochlorothiazide BP 25 mg	-do-	New	USFDA	<i>GB ḡīv c̄l̄q̄Rb b̄B m̄avq m̄m̄m-Gi 238 Zg mfviq Avt` b bugĀy K̄i evi m̄xvš- enj iLv h̄B̄tZ c̄t̄i /</i>	<i>GB ḡīv c̄l̄q̄Rb b̄B m̄avq Avt` b bugĀy K̄i nBj /</i>	
	y) <b>Amlodipine 5 mg + Valsartan 160 mg + Hydrochlorothiazide 12.50 mg Tablet</b>  Amlodipine Besilate BP 6.93 mg (eq. to Amlodipine 5 mg) + Valsartan USP 160 mg + Hydrochlorothiazide BP 12.5 mg	-do-	New	USFDA	<i>GB ḡīv c̄l̄q̄Rb b̄B m̄avq m̄m̄m-Gi 238 Zg mfviq Avt` b bugĀy K̄i evi m̄xvš- enj iLv h̄B̄tZ c̄t̄i /</i>	<i>GB ḡīv c̄l̄q̄Rb b̄B m̄avq Avt` b bugĀy K̄i nBj /</i>	
	z) <b>Amlodipine 5 mg + Valsartan 160 mg + Hydrochlorothiazide 25 mg Tablet</b>  Amlodipine Besilate BP 6.93 mg (eq. to Amlodipine 5 mg) + Valsartan USP 160 mg + Hydrochlorothiazide BP 25 mg	-do-	New	USFDA	<i>GB ḡīv c̄l̄q̄Rb b̄B m̄avq m̄m̄m-Gi 238 Zg mfviq Avt` b bugĀy K̄i evi m̄xvš- enj iLv h̄B̄tZ c̄t̄i /</i>	<i>GB ḡīv c̄l̄q̄Rb b̄B m̄avq Avt` b bugĀy K̄i nBj /</i>	
	aa) <b>Gliclazide 60 mg MR Tablet</b>  Gliclazide BP 60 mg	It is indicated for type II Diabetes in association with dietary measures when dietary measures alone are inadequate to control blood glucose.	30 mg & 80 mg Tablet		<i>GB ḡīv c̄l̄q̄Rb b̄B m̄avq m̄m̄m-Gi 238 Zg mfviq Avt` b bugĀy K̄i evi m̄xvš- enj iLv h̄B̄tZ c̄t̄i /</i>	<i>GB ḡīv c̄l̄q̄Rb b̄B m̄avq Avt` b bugĀy K̄i nBj /</i>	

<i>bs</i>	<i>cōZKvičKi bug</i>	<i>Jl̄ai bug I tR̄bniK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Aiče` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigilji 56 Zg mfvi m̄xviši</i>	<i>mfvi m̄xviš-</i>
16.	Nuvista Pharma Ltd.	a) <b>Estriol 2 mg Tablet</b>  Estriol USP 2 mg	Hormone replacement therapy for treatment of genito-urinary complaints due to estrogen deficiency in peri and post menopausal women. Infertility due to poor cervical penetration.	1 mg Tablet		<i>AvcizZt -MZ ivLv hvBtZ cv̄i Ges cōZōiblJtK cōqvRbq ti d̄ti Ymn cv̄ivq Aiče` b Kivi Rb ej v hvBtZ cv̄i /</i>	<i>AvcizZt -MZ ivLv nBj Ges cōZōiblJtK cōqvRbq ti d̄ti Ymn cv̄ivq Aiče` b Kivi Rb ej v nBj /</i>
17.	Beximco Pharmaceuticals Ltd.	a) <b>Vitamin A 0.63 mg + Vitamin C 66 mg + Vitamin D 12.5 mcg + Vitamin E 22 mg + Vitamin K 50 mcg + Riboflavin 0.400 mg + Calcium 100 mg Chewable Tablet</b>  Dry Vitamin A Acetate, Type 500-50 USP 3.700 mg (eq. to Vitamin A 0.63 mg) Ascorbic Acid (97% DC) USP 68.041 mg (eq. to Vitamin C 66 mg) Dry Vitamin D3, Colecalciferol 1,00,000 IU/g USP 5 mg (eq. to Vitamin D 12.5 mcg) Dry Vitamin E Acetate 50% USP 44 mg (eq. to Vitamin E 22 mg) Dry Vitamin K1 5% USP 1 mg (eq. to Vitamin K 50 mcg) Riboflavin USP 0.400 mg Calcium Carbonate (Heavy) BP 250 mg (eq. to Calcium 100 mg)	Indicated for total bone health and skin care.	New		<i>AvcizZt -MZ ivLv hvBtZ cv̄i /</i>	<i>AvcizZt -MZ ivLv nBj /</i>
		b) <b>Vitamin A 0.3 mg + Riboflavin 0.100 mg + Calcium Pantothenate 0.500 mg + Cyanocobalamin 1.0% 0.150 mg + Vitamin E 6 mg + Vitamin C 30 mg Chewable Tablet</b>  Dry Vitamin A Acetate, Type 500-50 USP 1.744 mg (eq. to Vitamin A 0.3 mg) Riboflavin USP 0.100 mg Calcium Pantothenate USP 0.500 mg Cyanocobalamine 1.0% USP 0.150 mg Dry Vitamin E Acetate 50% USP 12.00 mg eq. to Vitamin E 6 mg Ascorbic Acid, 97% DC USP 30.928 mg eq. to Vitamin C 30 mg	It is indicated to prevent aging.	New		<i>AvcizZt -MZ ivLv hvBtZ cv̄i /</i>	<i>AvcizZt -MZ ivLv nBj /</i>

<i>bs</i>	<i>cōZKvičKi bug</i>	<i>Jl̄ai bug I tRtbičK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Aiṭe` bKvix cōE</i> BNF or USFDA Reference	<i>tUKibK'ij mve-Kigilji 56 Zg mfvi mxiši</i>	<i>mfvi mxiš-</i>
	Beximco Pharmaceuticals Ltd.	<p>c) <b>Calcium 600 mg + Vitamin D 200 IU Chewable Tablet</b>  Calcium Carbonate USP 1500 mg (eq. to Calcium 600 mg) + Dry Vitamin D3, Colecalciferol 1,00,000 IU/G USP 2 mg (eq. to Vitamin D 200 IU)</p>	<ul style="list-style-type: none"> <li>Prevention and treatment of osteoporosis.</li> <li>For the treatment of hypocalcemic states.</li> <li>Dietary supplementation.</li> <li>Healthy bone formation and maintenance.</li> </ul>	500 mg + 200 IU		<i>AvcirZzt -MZ ivLv hvBtZ cvi i /</i>	<i>AvcirZzt -MZ ivLv nBj /</i>
		<p>d) <b>Distigmine Bromide 5 mg Tablet</b>  Distigmine Bromide INN 5 mg</p>	It is used in post operative urinary retention, prevention of ileus and intestinal atony following surgery, treatment of myasthenia gravis and treatment of neurogenic bladder.	New	BNF-57 Page-451	<i>Abtgr` b Kiv hvBtZ cvi i /</i>	<i>Abtgr` b Kiv nBj /</i>
		<p>e) <b>Sevelamer HCl 400 mg Tablet</b>  Sevelamer Hydrochloride INN 400 mg</p>	Indicated for the management of hyperphosphataemia in adult patients with stage 4 and 5 chronic renal failure on hemodialysis.	New	USFDA	<i>Abtgr` b Kiv hvBtZ cvi i /</i>	<i>Abtgr` b Kiv nBj /</i>
		<p>f) <b>Sevelamer HCl 800 mg Tablet</b>  Sevelamer Hydrochloride INN 800 mg</p>	-do-	New	USFDA	<i>Abtgr` b Kiv hvBtZ cvi i /</i>	<i>Abtgr` b Kiv nBj /</i>
		<p>g) <b>Fenoprofen 200 mg Capsule</b>  Fenoprofen Calcium USP 231.50 mg (eq. to Fenoprofen 200 mg)</p>	<ul style="list-style-type: none"> <li>For relief of mild to moderate pain in adults.</li> <li>For relief of the signs and symptoms of rheumatoid arthritis.</li> <li>For relief of the signs and symptoms of osteoarthritis.</li> </ul>	New	USFDA	<i>c̄qirRb bvB neaq Aiṭe` b bigAjy Kiv hvBtZ cvi i /</i>	<i>c̄qirRb bvB neaq Aiṭe` b bigAjy Kiv nBj /</i>

<i>bs</i>	<i>cōZKvi tKi bvg</i>	<i>Jl̄tai bvg I tR̄tbv K bvg</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-KvglJi 56 Zg mfvi m̄xvšl</i>	<i>mfvi m̄xvš-</i>
	Beximco Pharmaceuticals Ltd.	<p>h) <b>Fenoprofen 400 mg Capsule</b> Fenoprofen Calcium USP 463 mg (eq. to Fenoprofen 400 mg)</p> <p>i) <b>Fenoprofen 600 mg Tablet</b> Fenoprofen Calcium USP 694.50 mg (eq. to Fenoprofen 600 mg)</p> <p>j) <b>Racecadotril 100 mg Capsule</b> Racecadotril INN 100 mg</p> <p>k) <b>Simvastatin 20 mg + Niacin 750 mg Extended Release Tablet</b> Simvastatin USP 20 mg + Niacin USP 750 mg</p> <p>l) <b>Simvastatin 20 mg + Niacin 1000 mg Extended Release Tablet</b> Simvastatin USP 20 mg + Niacin USP 1000 mg</p> <p>m) <b>Olmesartan Medoxomil 40mg + Hydrochlorothiazide 12.5mg Tablet</b> Olmesartan Medoxomil INN 40mg + Hydrochlorothiazide BP 12.5mg</p> <p>n) <b>Olmesartan Medoxomil 40mg + Hydrochlorothiazide 25mg Tablet</b> Olmesartan Medoxomil INN 40mg + Hydrochlorothiazide BP 25mg</p> <p>o) <b>Amlodipine 5mg + Atorvastatin 20mg Tablet</b> Amlodipine Besilate 6.940mg eq. to Amlodipine 5mg + Atorvastatin Calcium 21.640mg eq. to</p>	<ul style="list-style-type: none"> <li>For relief of mild to moderate pain in adults.</li> <li>For relief of the signs and symptoms of rheumatoid arthritis.</li> <li>For relief of the signs and symptoms of osteoarthritis.</li> </ul> <p>-do-</p> <p>Acute diarrhea of presumed infectious origin and noninfectious origin.</p>	New  New  New  New  New	USFDA  USFDA  USFDA  USFDA  USFDA	<i>Abfgi` b Kiv hvBtZ cvti /</i> <i>cōqirb bvB neaq Avte` b bvgÄy Kiv hvBtZ cvti /</i> <i>cōqirb bvB neaq Avte` b bvgÄy Kiv hvBtZ cvti /</i> <i>GB ḡT v cōqirb bvB neaq m̄m̄m-Gi 238 Zg mfviq Avte` b bvgÄy Kvi evi m̄xvš- enj i Lv hvBtZ cvti /</i> <i>GB ḡT v cōqirb bvB neaq m̄m̄m-Gi 238 Zg mfviq Avte` b bvgÄy Kvi evi m̄xvš- enj i Lv hvBtZ cvti /</i> <i>Abfgi` b Kiv hvBtZ cvti /</i> <i>GB ḡT v cōqirb bvB neaq m̄m̄m-Gi 238 Zg mfviq Avte` b bvgÄy Kvi evi m̄xvš- enj i Lv hvBtZ cvti /</i> <i>GB ḡT v cōqirb bvB neaq m̄m̄m-Gi 238 Zg mfviq Avte` b bvgÄy Kvi evi m̄xvš- enj i Lv hvBtZ cvti /</i> <i>GB ḡT v cōqirb bvB neaq m̄m̄m-Gi 238 Zg mfviq Avte` b bvgÄy Kvi evi m̄xvš- enj i Lv hvBtZ cvti /</i>	<i>Abfgi` b Kiv nBj /</i> <i>cōqirb bvB neaq Avte` b bvgÄy Kiv nBj /</i> <i>cōqirb bvB neaq Avte` b bvgÄy Kiv nBj /</i> <i>GB ḡT v cōqirb bvB neaq Avte` b bvgÄy Kiv nBj /</i> <i>GB ḡT v cōqirb bvB neaq Avte` b bvgÄy Kiv nBj /</i> <i>Abfgi` b Kiv nBj /</i> <i>GB ḡT v cōqirb bvB neaq Avte` b bvgÄy Kvi evi m̄xvš- enj i Lv hvBtZ cvti /</i> <i>GB ḡT v cōqirb bvB neaq Avte` b bvgÄy Kvi evi m̄xvš- enj i Lv hvBtZ cvti /</i> <i>GB ḡT v cōqirb bvB neaq Avte` b bvgÄy Kvi evi m̄xvš- enj i Lv hvBtZ cvti /</i> <i>GB ḡT v cōqirb bvB neaq Avte` b bvgÄy Kvi evi m̄xvš- enj i Lv hvBtZ cvti /</i>

	Atorvastatin 20mg	cholesterol, LDL-Cholesterol.			<i>m×rš- enij i vLv hBtZ crti /</i>	
--	-------------------	-------------------------------	--	--	---	--

<i>bs</i>	<i>cőZKvičKi bug</i>	<i>Jl̄ai bug I tR̄buiK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt` bKvix cőE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kiugij 56 Zg mfvi m̄xviši</i>	<i>mfvi m̄xviš-</i>
	Beximco Pharmaceuticals Ltd.	p) <b>Amlodipine 5 mg + Hydrochlorothiazide 25 mg + Valsartan 160 mg Tablet</b>  Amlodipine Besilate BP 6.940 mg eq. to 5 mg Amlodipine + Hydrochlorothiazide BP 25 mg + Valsartan USP 160 mg	Treatment of hypertension.	New	USFDA	<i>GB ḡi v c̄qyRb bvB neaq m̄mim-Gi 238 Zg mfviq Avt` b bugAjy Kiv evi m̄xviš- enj i vLv hvBtZ cvi i /</i>	<i>GB ḡi v c̄qyRb bvB neaq Avt` b bugAjy Kiv nBj /</i>
	q) <b>Amlodipine 10 mg + Hydrochlorothiazide 12.50 mg + Valsartan 160 mg Tablet</b>  Amlodipine Besilate BP 13.880 mg eq. to 10 mg Amlodipine + Hydrochlorothiazide BP 12.50 mg + Valsartan USP 160 mg	Treatment of hypertension.	New	USFDA	<i>GB ḡi v c̄qyRb bvB neaq m̄mim-Gi 238 Zg mfviq Avt` b bugAjy Kiv evi m̄xviš- enj i vLv hvBtZ cvi i /</i>	<i>GB ḡi v c̄qyRb bvB neaq Avt` b bugAjy Kiv nBj /</i>	
	r) <b>Perindopril 3.395 mg + Amlodipine 5 mg Tablet</b>  Amlodipine Besylate BP 6.940 mg (eq. to Amlodipine 5 mg) + Perindopril Arginine INN 5.00 mg (eq. to Perindopril 3.396 mg)	Treatment of essential hypertension and or stable coronary artery disease, in patients already controlled with perindopril and amlodipine given currently at the same dose level.	New		<i>AvcZzt -MZ i vLv hvBtZ cvi i Ges c̄ZôvbiUjK c̄qyRbxq ti dñti Y mn c̄qyq Avt` b Kiv evi Rb ej v hvBtZ cvi i /</i>	<i>AvcZzt -MZ i vLv nBj Ges c̄ZôvbiUjK c̄qyRbxq ti dñti Y mn c̄qyq Avt` b Kiv evi Rb ej v nBj /</i>	
	s) <b>Perindopril 6.79 mg + Amlodipine 5 mg Tablet</b>  Amlodipine Besylate BP 6.940 mg (eq. to Amlodipine 5 mg) + Perindopril Arginine INN 10.00 mg (eq. to Perindopril 6.79 mg)	-do-	New		<i>c̄qyRb bvB neaq Avt` b bugAjy Kiv hvBtZ cvi i /</i>	<i>c̄qyRb bvB neaq Avt` b bugAjy Kiv nBj /</i>	
	t) <b>Trifluridine 1% Ophthalmic Solution</b>  Trifluridine INN 10 mg/ ml	Indicated for the treatment of primary keratoconjunctivitis and recurrent epithelial keratitis due to herpes simplex virus, types 1 and 2	1 mg & 5 mg Tablet	USFDA	<i>Abfgi` b Kiv hvBtZ cvi i /</i>	<i>Abfgi` b Kiv nBj /</i>	

<i>bs</i>	<i>cōZKvi tKi bug</i>	<i>Jl̄tai bug I tR̄tbuiK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Aute` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-KiugUj 56 Zg mfvi imxviši</i>	<i>mfvi imxviš-</i>
	Beximco Pharmaceuticals Ltd.	<p>u) <b>Moxifloxacin 0.5% + Dexamethasone 0.1% Ophthalmic Solution</b>  Moxifloxacin Hydrochloride INN 5.45 mg Eq. to Moxifloxacin 5 mg + Dexamethasone Sodium Phosphate USP 1.10 mg Eq. to Dexamethasone Phosphate 1 mg/ml</p> <p>Indicated for steroid responsive inflammatory ocular conditions for which a corticosteroid is indicated and where bacterial infection or a risk of bacterial ocular infections exists. The combination can also be used for post operative inflammation and any other ocular inflammation associated with infection.</p>	New			<i>AvcZZt -MZ ivLv hbZ cv̄ti Ges c̄ZōbiUtK c̄qivRbq ti d̄ti Y mn cv̄vq Aute` b Kvi evi Rb ej v hbZ cv̄ti /</i>	<i>AvcZZt -MZ ivLv nBj Ges c̄ZōbiUtK c̄qivRbq ti d̄ti Y mn cv̄vq Aute` b Kvi evi Rb ej v nBj /</i>
		<p>v) <b>Sodium Chloride 5% Ophthalmic Solution</b>  Sodium Chloride BP 50 mg/ ml</p> <p>It is used to reduce swelling of the cornea in certain eye conditions</p>	0.9 gm/100 ml Eye Drops			<i>Abfgv` b Kiv hbZ cv̄ti /</i>	<i>Abfgv` b Kiv nBj /</i>
		<p>w) <b>Bepotastine Besilate 1.5% w/v Ophthalmic Solution</b>  Bepotastine Besilate INN 15 mg/ ml</p> <p>It is indicated for the treatment of itching associated with signs and symptoms of allergic conjunctivitis.</p>	New	USFDA		<i>Abfgv` b Kiv hbZ cv̄ti /</i>	<i>Abfgv` b Kiv nBj /</i>
		<p>x) <b>Besifloxacin 0.6% Ophthalmic Suspension</b>  Besifloxacin Hydrochloride INN 6.63 mg (eq. to Besifloxacin 6 mg)/ml</p> <p>Indicated for the treatment of bacterial conjunctivitis.</p>	New	USFDA		<i>Abfgv` b Kiv hbZ cv̄ti /</i>	<i>Abfgv` b Kiv nBj /</i>
		<p>y) <b>Tramadol 37.5 mg + Paracetamol 500 mg Tablet</b>  Tramadol Hydrochloride BP 37.50 mg + Paracetamol BP 500 mg</p> <p>It is indicated for the short term treatment of mild to moderate acute pain.</p>	New			<i>Jlaiui Safety, Efficacy and Usefulness m̄ūtKgZvgZ c̄v̄bi Rb Aa'vcK KvRx `xb tḡm̄as , Xikv tḡm̄tKj Ktj R Ges Wlt Rb̄tq` kwdK, Rvcvb evsj t̄ k t̄dÜkic nmcvZij GubKU t̄dY Kiv hbZ cv̄ti /</i>	<i>Jlaiui Safety, Efficacy and Usefulness m̄ūtKgZvgZ c̄v̄bi Rb Aa'vcK KvRx `xb tḡm̄as , Xikv tḡm̄tKj Ktj R Ges Wlt Rb̄tq` kwdK, Rvcvb evsj t̄ k t̄dÜkic nmcvZij GubKU t̄dY Kiv hbZ cv̄ti /</i>

<i>bs</i>	<i>cōZKvičKi bvg</i>	<i>Jl̄tai bvg I tR̄tbvK bvg</i>	<i>ibt`Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt`bKvix cōE BNF or USFDA Reference</i>	<i>tUKibKvij mve-KvglUj 56 Zg mfvi m̄xvšl</i>	<i>mfvi m̄xvš-</i>
	Beximco Pharmaceuticals Ltd.	<p>z) <b>Tapentadol 50 mg Tablet</b> Tapentadol Hydrochloride INN 58.234 mg (eq. to 50 mg Tapentadol)</p> <p>aa) <b>Tapentadol 75 mg Tablet</b> Tapentadol Hydrochloride INN 87.351 mg (eq. to 75 mg Tapentadol)</p> <p>ab) <b>Tapentadol 100 mg Tablet</b> Tapentadol Hydrochloride INN 116.468 mg (eq. to 100 mg Tapentadol)</p> <p>ac) <b>Paracetamol 500 mg + Pamabrom 25 mg Tablet</b> Paracetamol BP 500 mg + Pamabrom USP 25 mg Tablet</p>	<p>It is used to treat moderate to severe pain</p> <p>-do-</p> <p>-do-</p> <p>It is indicated for the temporary relief of symptoms accompanying premenstrual syndrome such as cramps, headache, irritability, backache and mild to moderate aches and pains. To reduce the temporary excess water frequently associated with the premenstrual period it is most effective if taken at the first sign of discomfort, usually a few days before a period.</p>	<p>New</p> <p>New</p> <p>New</p> <p>New</p>	<p>USFDA</p> <p>USFDA</p> <p>USFDA</p> <p>USFDA KZR.Abtgmv`Z bq Ges BNF-G AŞ-fy` bq neaq Avte`b bigAý Kiv hbZ cüti  </p>	<p>Abtgv`b Kiv hbZ cüti  </p> <p>Abtgv`b Kiv hbZ cüti  </p> <p>Abtgv`b Kiv hbZ cüti  </p> <p>Abtgv`b Kiv nBj  </p>	<p>Abtgv`b Kiv nBj  </p> <p>Abtgv`b Kiv nBj  </p> <p>Abtgv`b Kiv nBj  </p> <p>USFDA KZR.Abtgmv`Z bq Ges BNF-G AŞ-fy` bq neaq Avte`b bigAý Kiv nBj  </p>
		<p>ad) <b>Cetirizine Hydrochloride 5 mg + Pseudoephedrine Hydrochloride 120 mg Extended Release Tabet</b> Cetirizine Hydrochloride BP 5 mg + Pseudoephedrine Hydrochloride BP 120 mg</p>	<p>It should be administered when both the antihistaminic properties of cetirizine hydrochloride and the nasal decongestant properties of pseudoephedrine Hydrochloride are desired. They are indicated for the relief of nasal and non-nasal symptoms associated with seasonal or perennial allergic rhinitis.</p>	<p>New</p>	<p>USFDA</p>	<p>c̄qıRb bıB neaq Avte`b bigAý Kiv hbZ cüti  </p>	<p>c̄qıRb bıB neaq Avte`b bigAý Kiv nBj  </p>

<i>bs</i>	<i>cōZKvičKi bug</i>	<i>Jl̄tai bug I tR̄tbuiK bug</i>	<i>ibt`Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt`bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigilji 56 Zg mfvi imxviši</i>	<i>mfvi imxviš-</i>
Beximco Pharmaceuticals Ltd.	ae) <b>Prasugrel 5 mg Tablet</b>  Prasugrel Hydrochloride INN 5.489 mg (eq. to Prasugrel 5 mg)	It is indicated to reduce the rate of thrombotic cardiovascular events in patients with acute coronary syndrome who are to be managed with percutaneous coronary intervention	New	USFDA	<i>Abtgi`b Kiv hibZ c̄ti  </i>	<i>Abtgi`b Kiv nBj  </i>	
	af) <b>Prasugrel 10 mg Tablet</b>  Prasugrel Hydrochloride INN 10.978 mg (eq. to Prasugrel 10 mg)	-do-	New	USFDA	<i>Abtgi`b Kiv hibZ c̄ti  </i>	<i>Abtgi`b Kiv nBj  </i>	
18. M/s. Ziska Pharmaceuticals Ltd.	ag) <b>Lymecycline 408 mg Capsule</b>  Lymecycline BP 408.00 mg (eq. to 300 mg Tetracycline)	Treatment of acne vulgaris, brucella & pasturella infections, pulmonary, GUT & ophthalmologic infecios due to Chlamydia, pulmonary & gut infectious due to mycoplasma, rickettsial infections, coxiella burnetti (Q fever), gonococcal infections, resp & bronchopulmonary infections due to H. influenzae, treponema, spirochetes, cholera	New	BNF-58 Page 309	<i>Abtgi`b Kiv hibZ c̄ti  </i>	<i>Abtgi`b Kiv nBj  </i>	
	a) <b>Famotidine 10 mg + Calcium Carbonate 800 mg + Magnesium Hydroxide 165 mg Tablet</b>  Famotidine BP 10 mg + Calcium Carbonate USP 800 mg + Magnesium Hydroxide BP 165 mg	The short term symptomatic relief of heartburn, indigestion, acid indigestion and hyperacidity.	New	USFDA	<i>Kifbkb Ac̄qibRbiq neaq Avt`b bigAij Kiv hibZ c̄ti  </i>	<i>Kifbkb Ac̄qibRbiq neaq Avt`b bigAij Kiv nBj  </i>	
	b) <b>Palonosetron 0.25 mg Tablet</b>  Palonosetron INN 0.25 mg	It is indicated for Chemotherapy induced Nausea and Vomiting	0.25mg/vial Injection		<i>USFDA KZR.Abtgi`Z bq Ges BNF-G Aš-fj bq neaq Avte`b bigAij Kiv hibZ c̄ti  </i>	<i>USFDA KZR.Abtgi`Z bq Ges BNF-G Aš-fj bq neaq Avte`b bigAij Kiv nBj  </i>	
	c) <b>Palonosetron 0.50 mg Tablet</b>  Palonosetron INN 0.50 mg	-do-	0.25 mg/Vial Injection		<i>USFDA KZR.Abtgi`Z bq Ges BNF-G Aš-fj bq neaq Avte`b bigAij Kiv hibZ c̄ti  </i>	<i>USFDA KZR.Abtgi`Z bq Ges BNF-G Aš-fj bq neaq Avte`b bigAij Kiv nBj  </i>	

<i>bs</i>	<i>cōZKvi tKi bug</i>	<i>Jl̄tai bug I tR̄tbuiK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Aute` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-KiugUj 56 Zg mfvi imxviši</i>	<i>mfvi imxviš-</i>
19.	The Acme Laboratories Ltd.	a) <b>Fluconazole 200 mg Tablet</b>  Fluconazole BP 200 mg	Indicated for the treatment of acute or recurrent vaginal candidiasis, mucosal candidiasis, systemic candidiasis and cryptococcal infections.	50 mg & 150 mg Tablet 50 mg, 150 mg & 200 mg Capsule		USFDA KZR Abtgw Z bq Ges BNF-G Aš-fq bq neaq Aute` b bugÄy Kiv hbZ cüti	USFDA KZR Abtgw Z bq Ges BNF-G Aš-fq bq neaq Aute` b bugÄy Kiv nBj
20.	Opso Saline Ltd.	a) <b>Sodium Chloride 50mg/ml Eye Drops</b>  Sodium Chloride BP 5 gm/100 ml	Indicated for the temporary relief of corneal edema, Bullous Keratopathy.	0.9 gm/100 ml Eye Drops		Abtgv` b Kiv hbZ cüti	Abtgv` b Kiv nBj
21.	Novartis Bangladesh Ltd.	a) <b>Tramadol HCl 37.5 mg + Paracetamol BP 325 mg Tablet</b>  Tramadol Hydrochloride BP 37.50 mg + Paracetamol BP 325 mg	It is indicated for the short term management of acute pain.	New	USFDA BNF-53 (Page 235)	<i>JlaiUi Safety, Efficacy and Usefulness māutK'gZvgZ cōvbi Rb Aa'vcK KvRx `xb tgimv-§, XvKv tgimvKj Ktj R Ges Wlt Rgytq` kwdK, Rvcv b evsj v k tdÜkic nwmcvZij GivbKU tcY Kiv hbZ cüti  </i>	<i>JlaiUi Safety, Efficacy and Usefulness māutK'gZvgZ cōvbi Rb Aa'vcK KvRx `xb tgimv-§, XvKv tgimvKj Ktj R Ges Wlt Rgytq` kwdK, Rvcv b evsj v k tdÜkic nwmcvZij GivbKU tcY Kiv nBj  </i>
		b) <b>Azithromycin 600 mg Tablet</b>  Azithromycin Dihydrate USP 628.84 mg (eq. to Azithromycin 600 mg)	Non gonococcal urethritis and cervicitis due to Chlamydia trachomatis, prophylaxis of disseminated mycobacterium avium complex (MAC) disease.	250 mg & 500 mg Tab. & Cap.	USFDA	<i>cōqibv bvb neaq Aute` b bugÄy Kiv hbZ cüti  </i>	<i>cōqibv bvb neaq Aute` b bugÄy Kiv nBj  </i>
		c) <b>Simvastatin 40 mg Tablet</b>  Simvastatin EP 40 mg	Treatment of hypercholesterolaemia and combined hyperlipidaemia	5mg, 10 mg & 20 mg Tab.	USFDA	<i>Abtgv` b Kiv hbZ cüti  </i>	<i>Abtgv` b Kiv nBj  </i>

<i>bs</i>	<i>cōZKvi tKi bvg</i>	<i>Jl̄tai bvg I tR̄tbwK bvg</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt̄e` bKvix cōĒ BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigij 56 Zg mfvi m̄x̄išl</i>	<i>mfvi m̄x̄iš-</i>
	Novartis Bangladesh Ltd.	d) <b>Fluconazole 100 mg Capsule</b>  Fluconazole BP 100 mg	Indicated for Vaginal candidiasis and candidal balanitis, Mucosal candidiasis including oropharyngeal, oesophageal, non-invasive bronchopulmonary infections, candiduria, mucocutaneous and chronic oral atrophic candidiasis	50mg, 150mg, 200 mg Capsule	USFDA	<i>cōqirb bvb neaq Avte`b bvgÄy Kiv hwbZ ci i  </i>	<i>cōqirb bvb neaq Avte`b bvgÄy Kiv nBj  </i>
		e) <b>Azithromycin 100 mg/5 ml Powder for Suspension</b>  Azithromycin USP 100 mg/5 ml	For the treatment of respiratory tract infections, skin and soft tissue infections, urinary tract and genitals infections.	200 mg/5 ml Suspension	USFDA	<i>cōqirb bvb neaq Avte`b bvgÄy Kiv hwbZ ci i  </i>	<i>cōqirb bvb neaq Avte`b bvgÄy Kiv nBj  </i>
		f) <b>Risperidone 3 mg Tablet</b>  Risperidone BP 3 mg	Indicated for the management of the manifestation of psychotic disorders	1mg, 2mg, & 4 mg Tab.	USFDA	<i>Abfgv`b Kiv hwbZ ci i  </i>	<i>Abfgv`b Kiv nBj  </i>
		g) <b>Mirtazapine 45 mg Tablet</b>  Mirtazepine BP 45 mg	Indicated for the treatment of depressive illness	15 mg & 30 mg Tab.	USFDA	<i>Abfgv`b Kiv hwbZ ci i  </i>	<i>Abfgv`b Kiv nBj  </i>
		h) <b>Lamotrigine 200 mg Dispersible Tablet</b>  Lamotrigine BP 200 mg	It is used as monotherapy and as an adjunct to treatment with other antiepileptics for partial seizures and primary and secondarily generalized tonic-clonic seizures.	25 mg, 50 mg & 100 mg Tablet	USFDA	<i>cōqirb bvb neaq Avte`b bvgÄy Kiv hwbZ ci i  </i>	<i>cōqirb bvb neaq Avte`b bvgÄy Kiv nBj  </i>
22.	Popular Infusions Ltd.	a) <b>Fluconazole 2mg/ml Infusion</b>  Fluconazole USP 200 mg/100 ml IV Infusion	Treatment and Prophylaxis of fungal infection.	50 mg & 150 mg Tab. & Cap. 50 mg/5ml Powder for Suspension	BNF-58 Page No. 335	<i>Abfgv`b Kiv hwbZ ci i  </i>	<i>Abfgv`b Kiv nBj  </i>

<i>bs</i>	<i>cōZKvičKi bvg</i>	<i>Jl̄tai bvg I tR̄tbvūK bvg</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigilji 56 Zg mfvi m̄xvši</i>	<i>mfvi m̄xvš-</i>
23.	Chemico Laboratories Ltd.	a) <b>Paracetamol 500 mg + Phenylephrine HCl 6.1 mg + Caffeine 25 mg Capsule</b>  Paracetamol BP 500 mg + Phenylephrine Hydrochloride BP 6.10 mg + Caffeine BP 25 mg	It is effective in relieving the symptoms associated with colds and flu, including relief of aches & pains, sore throats, headache, nasal congestion and lowering of temperature.	New		USFDA KZR.Abtgwm` Z bq Ges BNF-G Aš-fy bq neaq Avte`b bvgÄy Kiv hbZ c̄ti /	USFDA KZR.Abtgwm` Z bq Ges BNF-G Aš-fy bq neaq Avte`b bvgÄy Kiv nBj /
	b) <b>Magaldrate 800 mg Powder /Sachet</b>  Magaldrate USP 800 mg/Sachet	Indicated to relieve of symptoms of dyspepsia, heartburn, acid indigestion, sour stomach, gastro esophageal reflux disease and flatulence. It is also prescribed for hyperacidity with peptic ulcer, gastritis and esophagitis.	400 mg/5 ml Suspension & 400 mg Tablet		USFDA KZR.Abtgwm` Z bq Ges BNF-G Aš-fy bq neaq Avte`b bvgÄy Kiv hbZ c̄ti /	USFDA KZR.Abtgwm` Z bq Ges BNF-G Aš-fy bq neaq Avte`b bvgÄy Kiv nBj /	
	c) <b>Sucralfate 1 gm/10 ml Suspension</b>  Sucralfate USP 10 gm/100 ml	It is indicated in the short term treatment of active duodenal ulcer.	1 gm & 500 mg Tablet	USFDA	c̄qirb bq neaq Avte`b bvgÄy Kiv hbZ c̄ti /	c̄qirb bq neaq Avte`b bvgÄy Kiv nBj /	
	d) <b>Calcium 400 mg + Magnesium 150 mg + Zinc 5 mg + Copper 500 mcg + Manganese 0.25 mg + Selenium 25 mcg + Vitamin D 2.50 mcg + Boron 0.30 mg Tablet</b>  Calcium Carbonate USP 1000 mg (eq. to 400 mg Elemental Calcium) + Magnesium BP 150 mg + Zinc BP 5 mg + Copper BP 500 mcg + Manganese BP 0.25 mg + Selenium BP 25 mcg + Vitamin D BP 2.50 mcg + Boron BP 0.30 mg	Treatment of osteoporosis, osteomalacia, rickets, tetany and parathyroid disease.	New		AvcIZzt -MZ ivLv hbZ c̄ti /	AvcIZzt -MZ ivLv nBj /	

<i>bs</i>	<i>cōZKvičKi bvg</i>	<i>Jl̄tai bvg I tR̄tbvčK bvg</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibKvij mve-KvglUj 56 Zg mfvi m̄xvšl</i>	<i>mfvi m̄xvš-</i>
	Chemico Laboratories Ltd.	e) Cod Liver Oil 100 mg + Vitamin A 2000 IU + Vitamin D <sub>3</sub> 200 IU + Vitamin B <sub>1</sub> 0.70 mg, Vitamin B <sub>2</sub> 0.85 mg + Vitamin B <sub>6</sub> 0.35 mg + Vitamin C 17.50 mg + Vitamin E 1.50 mg + Nicotinamide 9.00 mg + Folic Acid 100 mcg + Vitamin B <sub>12</sub> 0.50 mcg + Biotin 75 mcg + Pantothenic Acid 3 mg/5 ml Syrup  Cod Liver Oil BP 2 gm + Vitamin A (as Vitamin A Propionate) BP 40,000 IU + Vitamin D <sub>3</sub> (as Cholecalciferol) BP 4000 IU + Vitamin B <sub>1</sub> (as Thiamin Hydrochloride) BP 14 mg + Vitamin B <sub>2</sub> (as Riboflavin Sodium Phosphate) BP 17 mg + Nicotinamide BP 180 mg + Vitamin B <sub>6</sub> (as Pyridoxine HCl) BP 7 mg + Vitamin E (as Alpha Tocopherol Acetate) BP 30 mg + Vitamin C (as Ascorbic Acid) BP 350 mg + Folic Acid BP 2000 mcg + Vitamin B <sub>12</sub> BP 10 mcg + Biotin BP 1500 mcg + Pantothenic Acid BP 60 mg/100 ml	To increase multivitamin to body for good health. Prevention of vitamin lacking symptom for growing child, lack of appetite child, including patient under rehabilitating period.	Vitamin A BP 40,000IU (as Vit. A Propionate) Vitamin D (as Colecalciferol) BP 4000IU Vitamin C (Ascorbic Acid) BP 0.35 gm Vitamin B1 (Thiamine HCl) BP 0.014 gm Vitamin B2 (Riboflavin Sodium Phosphate) BP 0.017 gm Vitamin B6 (Pyridoxine HCl) BP 0.007 gm Vitamin E (as Alpha Tocopherol) BP 0.030gm Nicotinamide BP 0.18gm Cod Liver Oil BP 2gm/100ml		AvcirZZt -MZ ivLv hvBtZ cvti /	AvcirZZt -MZ ivLv nBj /
24.	R.A.K. Pharmaceuticals Ltd.	a) <b>Voglibose 200 mcg Tablet</b>  Voglibose INN 200 mcg	Indicated for improvement of postprandial hyperglycemia in diabetes mellitus only when diet and/or exercise or oral hypoglycemic drug or insulin preparation in addition to diet and/or exercise does not result in adequate glycemic control.	New		USFDA KZR.Abtgmv Z bq / BNF-G Aš-fq bq wearq Avt` b bvgÄij Kiv hvBtZ cvti /	USFDA KZR.Abtgmv Z bq / BNF-G Aš-fq bq wearq Avt` b bvgÄij Kiv nBj /
		b) <b>Voglibose 300 mcg Tablet</b>  Voglibose INN 300 mcg	-do-	New		USFDA KZR.Abtgmv Z bq / BNF-G Aš-fq bq wearq Avt` b bvgÄij Kiv hvBtZ cvti /	USFDA KZR.Abtgmv Z bq / BNF-G Aš-fq bq wearq Avt` b bvgÄij Kiv nBj /

<i>bs</i>	<i>cōZKvi tKi bvg</i>	<i>Jl̄tai bvg I tR̄tbv K bvg</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avtē` bKvix cōĒ BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigij 56 Zg mfvi imxvši</i>	<i>mfvi imxvš-</i>
R.A.K. Pharmaceuticals Ltd.	c)	<b>Diclofenac Sodium 1.5% Topical Solution</b>  Diclofenac Sodium USP 1.50 gm/100 ml	It is indicated for treatment of the symptoms associated with osteoarthritis of the knee only, for a treatment regimen of not more than three months duration, whether continuous or intermittent.	100 mg Tab. & Cap. 1 gm/100 gm Gel 12.50 mg & 60 mg Suppository	USFDA	<i>cōqirb bvB neaq Avte`b bvgāy Kiv hBtZ cti  </i>	<i>cōqirb bvB neaq Avte`b bvgāy Kiv nBj  </i>
	d)	<b>Flecainide Acetate 100 mg Tablet</b>  Flecainide Acetate USP 100 mg	Indicated for the prevention of Paroxysmal supraventricular tachycardias, including atrioventricular nodal reentrant tachycardia, atrioventricular reentrant tachycardia and other supraventricular tachycardias of unspecified mechanism associated with disabling symptoms.	New	USFSA	<i>Abtgy`b Kiv hBtZ cti  </i>	<i>Abtgy`b Kiv nBj  </i>
	e)	<b>Ciprofloxacin 100 mg Tablet</b>  Ciprofloxacin Hydrochloride USP 111.00 mg (eq. to Ciprofloxacin 100 mg)	It is used to treat kidney or bladder (urinary tract) infections.	250 mg & 500 mg Tab.	BNF-56 (Page 321)	<i>cōqirb bvB neaq Avte`b bvgāy Kiv hBtZ cti  </i>	<i>cōqirb bvB neaq Avte`b bvgāy Kiv nBj  </i>
	f)	<b>Amlodipine 10 mg + Telmisartan 40 mg Tablet</b>  Amlodipine Besylate BP 13.80 mg (eq. to 10 mg Amlodipine) + Telmisartan INN 40 mg	Indicated for the treatment of hypertension, alone or with other antihypertensive agents. It may be used as initial therapy in patients who are likely to need multiple drugs to achieve their blood pressure goals.	New	USFDA	<i>cōqirb bvB neaq Avte`b bvgāy Kiv hBtZ cti  </i>	<i>cōqirb bvB neaq Avte`b bvgāy Kiv nBj  </i>

<i>bs</i>	<i>cōZKvi tKi bug</i>	<i>Jl̄tai bug I tR̄tbuiK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kiguij 56 Zg mfvi imxviši</i>	<i>mfvi imxviš-</i>
	R.A.K. Pharmaceuticals Ltd.	<p>g) <b>Amlodipine 10 mg + Telmisartan 80 mg Tablet</b> Amlodipine Besylate BP 13.80 mg (eq. to 10 mg Amlodipine) + Telmisartan INN 80 mg</p> <p>h) <b>Amlodipine 5 mg + Telmisartan 40 mg Tablet</b> Amlodipine Besylate BP 6.90 mg (eq. to 5 mg Amlodipine) + Telmisartan INN 40 mg</p> <p>i) <b>Amlodipine 5 mg + Telmisartan 80 mg Tablet</b> Amlodipine Besylate BP 6.90 mg (eq. to 5 mg Amlodipine) + Telmisartan INN 80 mg</p> <p>j) <b>Aliskiren 150 mg + Valsartan 160 mg Tablet</b> Aliskiren INN 150 mg + Valsartan USP 160 mg</p> <p>k) <b>Aliskiren 300 mg + Valsartan 320 mg Tablet</b> Aliskiren INN 300 mg + Valsartan USP 320 mg</p> <p>l) <b>Vitamin A 6600 IU + Vitamin C 400 mg + Vitamin E 150 IU + Riboflavin 10 mg + Zinc 60 mg + Copper 4 mg + Manganese 10 mg + Selenium 40 mcg + Lutein/Zeaxanthin 4 mg Tablet</b> Vitamin A BP 6600 IU + Vitamin C BP 400 mg + Vitamin E USP 150 IU + Riboflavin BP 10 mg + Zinc (as Zinc Acetate) BP 60 mg + Manganese BP 10 mg + Copper USP 4 mg + Selenium BP 40 mcg + Lutein/Zeaxanthin USP 4 mg</p>	<p>Indicated for the treatment of hypertension, alone or with other antihypertensive agents. It may be used as initial therapy in patients who are likely to need multiple drugs to achieve their blood pressure goals.</p> <p>-do-</p> <p>-do-</p> <p>It is indicated for the treatment of hypertension.</p> <p>-do-</p> <p>Indicated for age related eye disease e.g. age related muscular degeneration or people with frequent exposure to sunlight.</p>	<p>New</p> <p>New</p> <p>New</p> <p>New</p> <p>New</p>	<p>USFDA</p> <p>USFDA</p> <p>USFDA</p> <p>USFDA</p> <p>USFDA</p>	<p>cōqiqRb bvB neaq Avte`b bigÄy Kiv hvBtZ cvti  </p> <p>Abfgi`b Kiv hvBtZ cvti  </p> <p>cōqiqRb bvB neaq Avte`b bigÄy Kiv hvBtZ cvti  </p> <p>cōqiqRb bvB neaq Avte`b bigÄy Kiv hvBtZ cvti  </p> <p>cōqiqRb bvB neaq Avte`b bigÄy Kiv hvBtZ cvti  </p> <p>AvciZZt -MZ ivLv hvBtZ cvti  </p>	<p>cōqiqRb bvB neaq Avte`b bigÄy Kiv nBj  </p> <p>Abfgi`b Kiv nBj  </p> <p>cōqiqRb bvB neaq Avte`b bigÄy Kiv nBj  </p> <p>cōqiqRb bvB neaq Avte`b bigÄy Kiv nBj  </p> <p>cōqiqRb bvB neaq Avte`b bigÄy Kiv nBj  </p> <p>AvciZZt -MZ ivLv nBj  </p>
25.	Jayson Pharmaceuticals Ltd.	<p>a) <b>Lidocaine Hydrochloride 2 gm + Epinephrine 1.25 mg/100 ml Dental cartridge</b> Lidocaine Hydrochloride USP 2.133 gm (eq. to Anhydrous Lidocaine Hydrochloride 2 gm) + Ephinephrine Bitartrate USP 2.274 mg (eq. to Epinephrine (Adrenaline) 1.250 mg)/100 ml</p>	<p>It is a local anesthetic used in dental infiltration anaesthesia and all dental nerve block techniques.</p>	<p>New</p>	<p>BNF-55 (strength উলঞ্চা biB)</p>	<p>Abfgi`b Kiv hvBtZ cvti  </p>	<p>Abfgi`b Kiv nBj  </p>

<i>bs</i>	<i>cōZKvičKi bug</i>	<i>Jl̄ai bug I tR̄buiK bug</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Aute` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigilji 56 Zg mfvi imxviši</i>	<i>mfvi imxviš-</i>
26.	General Pharmaceuticals Ltd.	<p>a) <b>Montelukast 10 mg + Levocetirizine Dihydrochloride 5 mg Tablet</b>  Montelukast Sodium INN 10.40 mg (eq. to 10 mg Montelukast) + Levocetirizine Dihydrochloride INN 5 mg</p> <p>b) <b>Montelukast 10 mg + Bambuterol Hydrochloride 10 mg Tablet</b>  Montelukast Sodium INN 10.400 mg (eq. to 10 mg Montelukast) + Bambuterol Hydrochloride BP 10 mg</p> <p>c) <b>Montelukast 4 mg + Bambuterol Hydrochloride 5 mg Tablet</b>  Montelukast Sodium INN 4.160 mg (eq. to 4 mg Montelukast) + Bambuterol Hydrochloride BP 5 mg</p>	<p>It is indicated for relief and/or prevention of symptoms of allergic rhinitis (seasonal and perennial), long term treatment of asthma and chronic urticaria.</p> <p>It is indicated for chronic treatment of asthma, bronchospasm and/or reversible airways obstruction in patients 12 years of age and older.</p> <p>It is indicated for chronic treatment of asthma, bronchospasm and/or reversible airways obstruction in patients 2-12 years of age.</p>	New		USFDA KZR.Abtgwi Z bq / BNF-G Aš-fq bq næarq Aute` b bugÄy Kiv hibtz cüti /	USFDA KZR.Abtgwi Z bq / BNF-G Aš-fq bq næarq Aute` b bugÄy Kiv nBj /
27.	Reman Drug Laboratories Ltd.	<p>a) <b>Norfloxacin Eye/Ear Drops 0.3%</b>  Norfloxacin USP 0.3 gm/100 ml</p>	<p>It is a fluoroquinolone carboxylic acid antibacterial agent for ocular-otic administration. Norfloxacin inhibits bacterial deoxyribonucleic acid synthesis and is bactericidal Eye : This drop is indicated for the treatment of the superficial infections of the eye and its annex: Conjunctivitis, blepharitis, tarsitis caused by pathogenic bacteria sensitive to norfloxacin Ear : It is indicated in severe and long lasting bacterial infection in otitis media, otitis externa It is also used in otorrhoea and infection caused by P. Vulgaris, Pseudomonas Aeruginosa, which are resistant to many other antibiotics.</p>	New	USFDA	Abtgv` b Kiv hibtz cüti /	Abtgv` b Kiv nBj /

<i>bs</i>	<i>cōZKvi tKi bvg</i>	<i>Jl̄tai bvg I tR̄tbwK bvg</i>	<i>ibt` Rbv</i>	<i>Status (New Molecule/ Existing)</i>	<i>Avt` bKvix cōE BNF or USFDA Reference</i>	<i>tUKibK'ij mve-Kigij 56 Zg mfvi m̄xvšl</i>	<i>mfvi m̄xvš-</i>
28.	Eskayef Bangladesh Ltd.	a) <b>Prasugrel 5 mg Tablet</b>  Prasugrel Hydrochloride INN 5.49 mg (Eq. to Prasugrel 5 mg)	Indicated for the prevention of atherothrombotic events in patients with acute coronary syndrome.	New		<i>Abfgi` b Kiv h̄BtZ c̄ti  </i>	<i>Abfgi` b Kiv nBj  </i>
		b) <b>Prasugrel 10 mg Tablet</b>  Prasugrel Hydrochloride INN 10.98 mg (Eq. to Prasugrel 10 mg)	-do-	New		<i>Abfgi` b Kiv h̄BtZ c̄ti  </i>	<i>Abfgi` b Kiv nBj  </i>
29.	Techno Drugs	a) <b>Oxaliplatin 100mg/Vial Injection (Lyophilized)</b>  Oxaliplatin BP 100 mg/Vial	Treatment of metastatic colorectal cancer in combination with fluorouracil and folinic acid, colon cancer.	50mg/vial	BNF-57 Page 479	<i>Abfgi` b Kiv h̄BtZ c̄ti  </i>	<i>Abfgi` b Kiv nBj  </i>
30.	Renata Ltd.	a) <b>Levomenthol 35.55mg+ Thymol 3.15mg+ Chlorobutanol 2.25mg+ Terpineol 66.6mg+ Pumilio Pine oil 103.05mg+ Pine oil Sylvestris 9.00mg Soft gelatin Capsule</b>  Levomenthol BP 35.55mg+ Thymol BP 3.15mg+ Chlorobutanol BP 2.25mg+Terpineol BP 66.6mg+ Pumilio Pine oil Ph.Grade 103.05mg+ Pine oil Sylvestris BP 9.00mg	Indicated for symptomatic relief in nasal congestion, coryza and bronchitis	New		<i>c̄qyRb bvB neaq m̄m̄m-Gi 238 Zg mfvi Avt` b bvgÄy Kii evi m̄xvš- enij ivLv h̄BtZ c̄ti  </i>	<i>c̄qyRb bvB neaq Avt` b bvgÄy Kiv nBj  </i>
		b) <b>Salmon Oil Soft gelatin Capsule</b>  Salmon Oil BP 1000 mg (eq. to EPA 80 mg & DHA 120 mg)	Omega-3 fatty acids, also known as fish oils, have been used to treat rheumatoid arthritis, high blood pressure, high lipid (fat) levels in the blood, certain skin/scalp conditions, ulcerative colitis, or certain blood circulation problems.	New		<i>USFDA KZR Abfgi` Z bq Ges BNF-G Aš-fy bq neaq Avt` b bvgÄy Kiv h̄BtZ c̄ti  </i>	<i>USFDA KZR Abfgi` Z bq Ges BNF-G Aš-fy bq neaq Avt` b bvgÄy Kiv nBj  </i>

2.2 WMM KtUij Kiguij 238 Zg mfvi mxiš-i cmitcmtZ wbtgewYZ c` wj i Safety mxišKZ Z w`/idvti Y cñZövb KZR. `wLj Kiv nBqitQ t

bs	cñZKvičKi bug	Jltači bug I tRtbwK bug	ibt`Rbv	Status (New Molecule/ Existing)	Arte`bKvix cñE BNF or USFDA Reference	tUKubKij mve-Kiguij 56 Zg mfvi mxiši	mfvi mxiš-
01.	Opso Saline Ltd.	a) <b>Moxifloxacin160mg/100ml IV infusion</b>  Moxifloxacin Hydrochloride BP 174.56mg eq. to Moxifloxacin 160mg/100ml	Acute Bacterial sinusitis, acute bacterial exacerbation of chronic bronchitis, community acquired Pneumonia.	5 mg/ml Eye Drops & 400 mg Tab.	USFDA	Abtgr`b Kiv hBtZ cüti	Abtgr`b Kiv hBtZ cüti
02.	ACI Ltd.	a) <b>Cefditoren 200 mg Tablet</b>  Cefditoren Pivoxil INN 245 mg (eq. to 200 mg Cefditoren)	Uncomplicated skin & skin structure infections, Community-Acquired Pneumonia, Acute Bacterial Exacerbation of chronic Bronchitis, Pharyngitis/ Tonsillitis.		USFDA	Abtgr`b Kiv hBtZ cüti	Abtgr`b Kiv hBtZ cüti
		b) <b>Cefditoren 400 mg Tablet</b>  Cefditoren Pivoxil INN 490 mg (eq to 400 mg Cefditoren)	Uncomplicated skin & skin structure infections, Community-Acquired Pneumonia, Acute Bacterial Exacerbation of chronic Bronchitis, Pharyngitis/ Tonsillitis.			Abtgr`b Kiv hBtZ cüti	Abtgr`b Kiv hBtZ cüti
03.	Incepta Pharmaceuticals Ltd.	a) <b>Paracetamol 10mg/ml IV Infusion</b>  Paracetamol BP 1 gm/100 ml	For the relief of mild to moderate pain and the reduction of fever where an intravenous route of administration is considered clinically necessary.	500 mg Tablet & 120 mg/5 ml Suspension	BNF-57, Page 232	JlaiUi Safety, Efficacy and Usefulness mxišKigZ cñvbi Rb` Aa'icK KivRi `ib tgnir=§, XivKv tgwtKj Ktj R, Aa'icK gvi,e tnitmb mriRi x nefIM, XivKv tgwtKj Ktj R Ges Wt Rytq` kndK, Rvcib eisj it`k tdÜkic nmcizij GiubKU tcY Kiv hBtZ cüti	JlaiUi Safety, Efficacy and Usefulness mxišKigZ cñvbi Rb` Aa'icK KivRi `ib tgnir=§, XivKv tgwtKj Ktj R, Aa'icK gvi,e tnitmb mriRi x nefIM, XivKv tgwtKj Ktj R Ges Wt Rytq` kndK, Rvcib eisj it`k tdÜkic nmcizij GiubKU tcY Kiv hBtZ cüti
04.	Beximco Pharmaceuticals Ltd.	a) <b>Doxofylline 400 mg Tablet</b>  Doxofylline INN 400 mg	Reversible airways obstruction.			cñqirb bvB neaq Arte`b bvgAy Kiv hBtZ cüti	cñqirb bvB neaq Arte`b bvgAy Kiv hBtZ cüti

## 2.3 Proposed Product for Locally manufacture (Veterinary)

<i>bs</i>	<i>cōZKvičKi bvg</i>	<i>Jl̄tai bvg I tRtbi K bvg</i>	<i>ibt`Rbv</i>	<i>tUKibKvij mve-Kigui 56 Zg mfvi m̄xvši</i>	<i>mfvi m̄xvš-</i>
01.	The Acme Laboratories Ltd.	a) <b>Ivermectin &amp; Clorsulon Injection (Vet)</b>  Ivermectin BP 1 gm + Clorsulon USP 10 gm/100 ml	Indicated for the effective treatment and control of the Gastrointestinal roundworms, Lungworms and Liver flukes of cattle.	<i>c̄qvRb bvB meaq wlm̄m-Gi 238 Zg mfvq Avte` b bvgAij Kii evi m̄xvš- enij ivlv h̄BtZ c̄ti /</i>	<i>c̄iU tUKibKvij mve-Kigui cieZi mfvi c̄jivq Dc̄v̄bi Rb̄ ej v nBj /</i>
		b) <b>Amoxicillin 1 gm Bolus (Vet)</b>  Amoxicillin Trihydrate BP 1.150 gm (eq. to Amoxicillin 1gm)	It is indicated for the treatment of infections caused by microorganisms sensitive to amoxicillin in cattle, buffalo, sheep, goat, dog and cat.	<i>Ab̄gr`b Kiv h̄BtZ c̄ti /</i>	<i>Ab̄gr`b Kiv h̄BtZ c̄ti /</i>

## 2.4 Proposed Product for Import (Human)

bs	cūZKvitKi bvg	Jl t̄ai bvg I tRt̄bwi K bvg	mbt̄ Rbv	CPP/FSC	tUKibKij mve-KiguiJi 56 Zg mfvi m̄xvši	mfvi m̄xvš-
01.	Orion Corporation, Finland  (Novartis Bangladesh Ltd.)	a) <b>Stalevo 75 F.C Tablet</b>  Levodopa USP 75 mg + Carbidopa USP 18.75 mg + Entacapone INN 200 mg	Treatment of patients with parkinson's disease either at initiation of levodopa therapy or when end of dose motor fluctuations are not stabilized on levodopa/dopa decarboxylase inhibitor treatment.	EMEA	cūqRb bVB neaq Arte`b bigĀj Kiv hVbtZ c̄ti	cūqRb bVB neaq Arte`b bigĀj Kiv nBj
		b) <b>Stalevo 125 F.C Tablet</b>  Levodopa USP 125 mg + Carbidopa USP 31.25 mg + Entacapone INN 200 mg	-do-	EMEA	cūqRb bVB neaq Arte`b bigĀj Kiv hVbtZ c̄ti	cūqRb bVB neaq Arte`b bigĀj Kiv nBj
02.	S.A Alcon-Couvreur n.v., Belgium  (Globex Marketing Co. Ltd.)	a) <b>DUOVISC -Viscoelastic System</b>  <u>Provisc</u> : Sodium Hyaluronate 10 mg/ml <u>Viscoat</u> : Sodium Chondroitin Sulfate 40 mg + Sodium Hyaluronate 30 mg/ml	Indicated for use as an ophthalmic surgical aid in the anterior segment during cataract extraction and intraocular lens implantation.	Belgium	cūqRb bVB neaq Arte`b bigĀj Kiv hVbtZ c̄ti	cūqRb bVB neaq Arte`b bigĀj Kiv nBj
03.	Novartis Vaccines and Diagnostics GmbH & Co. KG, Germany  (Novartis Bangladesh Ltd.)	a) <b>Tetanol pur Suspension for Injection in Prefilled Syringe (Adsorbed Titanus Vaccine)</b>  Purified Tetanus Toxoid Concentrate not less than 40 IU/0.5 ml	Indicated for the active immunization against tetanus for infants aged 2 months and over, children, adolescents and adults.	Germany	cūqRb bVB neaq Arte`b bigĀj Kiv hVbtZ c̄ti	cūqRb bVB neaq Arte`b bigĀj Kiv nBj
		b) <b>Td-pur Suspension for Injection in Prefilled Syringe</b>  Adsorbed Diphtheria Toxoid not less than 2 IU + Adsorbed Tetanus Toxoid Not less than 20 IU/0.5 ml	Indicated for the active immunization against tetanus and diphtheria in individuals aged 5 years old and over.	Germany	cūqRb bVB neaq Arte`b bigĀj Kiv hVbtZ c̄ti	cūqRb bVB neaq Arte`b bigĀj Kiv nBj

<b>bs</b>	<b>cÜZKvičKi big</b>	<b>Jl̄tai big I tRtbwíK big</b>	<b>ibt`Rbv</b>	<b>CPP/FSC</b>	<b>tUKibKij mve-KigijIi 56 Zg mfvi wmxvš/</b>	<b>mfvi wmxvš-</b>
04.	Novartis Pharma AG, Switzerland  (Novartis Bangladesh Ltd.)	a) <b>Simulect 20 mg Powder for Solution for Injection (Lyophilisate for Injection)</b>  Basiliximab 20 mg/vial	Indicated for the prophylaxis of acute organ rejection in de novo renal transplantation in adult and paediatric patients.	Switzerland.	<i>Abfgv`b Kiv hBtZ cvti /</i>	<i>Abfgv`b Kiv nBj /</i>
		b) <b>Illaris 150 mg/ml Powder for Solution for Injection</b>  Canakinumab 150 mg/ml	Treatment of Cryopyrin Associated periodic syndromes in adults and children aged 4 years and older.	Switzerland.	<i>c̄qRb bvB neavq Arte`b bigÄy Kiv hBtZ cvti /</i>	<i>c̄qRb bvB neavq Arte`b bigÄy Kiv nBj /</i>
05.	Sanofi Pasteur S.A., France  (Sanofi-Aventis Bd. Ltd.)	a) <b>Stamaril Vaccine (Powder and solvent for suspension for injection in prefilled syringe)</b>  Yellow fever virus 17D-204 strain (live attenuated) Ph. Eur ≥1000 LD <sub>50</sub> units / 1 dose (0.5ml)	Indicated for active immunization against yellow fever.	France	<i>Abfgv`b Kiv hBtZ cvti /</i>	<i>Abfgv`b Kiv nBj /</i>
		b) <b>Tetrarixim Vaccine ( Suspension for injection in prefilled syringe)</b>  Diphtheria toxoid ≥ 30 IU Tetanus Toxoid ≥ 40 IU Bordetella pertussis antigens: Toxoid 25 µg Filamentous haemagglutinin 25 µg -Inactivated type 1 poliomyelitis virus 40 DU -Inactivated type 2 poliomyelitis virus 8 DU -Inactivated type 3 poliomyelitis virus 32 DU / 1 dose (0.5ml)	Indicated in the joint prevention of diphtheria, tetanus, pertussis and poliomyelitis.	France	<i>Abfgv`b Kiv hBtZ cvti /</i>	<i>Abfgv`b Kiv nBj /</i>

<b>bs</b>	<b>cőZKvičKi bvg</b>	<b>Jl t̄ai bvg I tRtbwi K bvg</b>	<b>ibt`Rbv</b>	<b>CPP/FSC</b>	<b>tUKibKij mve-KugijIi 56 Zg mfvi wmxvš/</b>	<b>mfvi mxvš-</b>
	Sanofi Pasteur S.A., France (Sanofe-Aventis Bd. Ltd.)	c) <b>VAXIGRIP Vaccine ( Suspension for injection in prefilled syringe)</b>  A/Brisbane/59/2007(H1N1)- like strain (A/Brisbane/59/2007 (IVR-148) 15µg HA  A/Brisbane/10/2007(H3N2)- like strain (A/Uruguay/716/2007 (NYMC X -175C) 15µg HA  B/Brisbane/60/2008- like strain B/Brisbane/60/2008 15µg HA / 1 dose (0.5ml)	The vaccine is indicated for the prevention of Influenza infection.	France	Abfgv`b Kiv hBtZ cv̄i	Abfgv`b Kiv nBj
06.	Les Laboratories Servier Industrie, France (Transcom Distribution Co., Ltd.)	a) <b>Coveram 5 mg/5 mg Tablet</b>  Perindopril arginine 5 mg eq. to Perindopril 3.395 mg + Amlodipine besilate Eur. Ph. 6.935 mg eq. to Amlodipine 5 mg  b) <b>Coveram 5 mg/10 mg Tablet</b>  Perindopril arginine 5 mg eq. to Perindopril 3.395 mg + Amlodipine besilate Eur. Ph. 13.870 mg eq. to Amlodipine 10 mg	Essential hypertension, Stable coronary artery disease, stroke.  -do-	France	c̄qRb bvB neaq Avte`b bvgÄj Kiv hBtZ cv̄i	c̄qRb bvB neaq Avte`b bvgÄj Kiv nBj
07.	Novo Nordisk A/S, Denmark (Transcom Distribution Ltd.)	a) <b>Victoza 6 mg/ml Solution for Injection in pre-filled pen</b>  <u>Quantity per unit dose Contains :</u> Liraglutide 6.0 mg/ml (Liraglutide – a Human glucagon-like peptide-1 (GLP-1) analogue produced by recombinant DNA technology)	It is used for the treatment of type 2 diabetes mellitus.	Denmark EMEA	Abfgv`b Kiv hBtZ cv̄i	Abfgv`b Kiv nBj
08.	Gland Pharma Ltd. India	a) <b>Paromomycin 375 mg/ml, 2 ml Ampoule for IM Use only</b>  Paromomycin 750 mg/2 ml	Indicated for intestinal amebiasis acute and chronic.	India	RiZiq Jla bmZ 2005 tḡZitēK Dbz 7iU t̄t̄ki tKvbiUj GdGmim `wLj Kii tZ cv̄i bvB neaq Avte`b bvgÄj Kiv hBtZ cv̄i	RiZiq Jla bmZ 2005 tḡZitēK Dbz 7iU t̄t̄ki tKvbiUj GdGmim `wLj Kii tZ cv̄i bvB neaq Avte`b bvgÄj Kiv nBj

<b>bs</b>	<b>cőZKvitKi big</b>	<b>Jl̄tai big I tR̄tbwi K big</b>	<b>ibt̄ Rbv</b>	<b>CPP/FSC</b>	<b>tUKibKij mve-KugijIi 56 Zg mfvi wmxvši</b>	<b>mfvi wmxvš-</b>
09.	H. Lundbeck A/S, Denmark (Lilac Pvt. Ltd.)	a) <b>Serdolect Film Coated Tablet 4 mg</b>  Sertindole 4 mg	It is indicated for the treatment of Schizophrenia.	Denmark UK	<i>Abfgv`b Kiv hBtZ c̄ti  </i>	<i>Abfgv`b Kiv nBj  </i>
		b) <b>Serdolect Film Coated Tablet 12 mg</b>  Sertindole 12 mg	-do-	Denmark UK	<i>Abfgv`b Kiv hBtZ c̄ti  </i>	<i>Abfgv`b Kiv nBj  </i>
		c) <b>Serdolect Film Coated Tablet 16 mg</b>  Sertindole 16 mg	-do-	Denmark UK	<i>Abfgv`b Kiv hBtZ c̄ti  </i>	<i>Abfgv`b Kiv nBj  </i>
		d) <b>Serdolect Film Coated Tablet 20 mg</b>  Sertindole 20 mg	-do-	Denmark UK	<i>Abfgv`b Kiv hBtZ c̄ti  </i>	<i>Abfgv`b Kiv nBj  </i>
10.	Pfizer Italia Srl, Italy (Janata Traders)	a) <b>Medrol 4 mg Tablet</b>  Methyl Prednisolone USP 4 mg	Indicated for conditions requiring glucocorticoids activity such as Endocrine Rheumatic collagen disorders/arthritis, dermatological allergic states, ophthalmic, disorder.	Italy UK	<i>Abgv`b Kiv hBtZ c̄ti  </i>	<i>Abgv`b Kiv nBj  </i>
		b) <b>Medrol 16 mg Tablet</b>  Methyl Prednisolone USP 16 mg	-do-	Italy UK	<i>Abfgv`b Kiv hBtZ c̄ti  </i>	<i>Abfgv`b Kiv nBj  </i>
10.	Pfizer Manufacturing Deutschland, Germany (Janata Traders)	a) <b>Champix 0.5 mg F/C Tablet</b>  Varenicline Tartrate 0.5 mg	Indicated for smoking cessation in adults.	European Union	<i>Abfgv`b Kiv hBtZ c̄ti  </i>	<i>Abfgv`b Kiv nBj  </i>
		b) <b>Champix 1 mg F/C Tablet</b>  Varenicline Tartrate 1 mg	-do-	European Union	<i>Abfgv`b Kiv hBtZ c̄ti  </i>	<i>Abfgv`b Kiv nBj  </i>
11.	Pfizer SA, Belgium (Janata Traders)	a) <b>Sayana 104 mg/0.65 ml Suspension for Injection</b>  Medroxyprogesterone Acetate USP 104 mg/ 0.65 ml	Indicated for long term female contraception.	Belgium UK	<i>c̄qRb b̄B weaq Āte`b bigÄj Kiv hBtZ c̄ti  </i>	<i>c̄qRb b̄B weaq Āte`b bigÄj Kiv nBj  </i>

<b>bs</b>	<b>cūZKvičKi big</b>	<b>Jl̄tai big I tRt̄bwi K big</b>	<b>ibt̄ Rbv</b>	<b>CPP/FSC</b>	<b>tUKibKij mve-KigijIi 56 Zg mfvi m̄xvši</b>	<b>mfvi m̄xvš-</b>
12.	Pfizer Limited, India  (Janata Traders)	a) <b>Magnex 1 gm Vial</b>  Sulbactam Sodium USP eq. to Sulbactam 500 mg + Cefoperazone Sodium IP eq. to Cefoperazone 500 mg/Vial	Indicated for the treatment of Respiratory tract infections, Urinary tract infections, peritonitis, cholecistitis, cholangitis and other intra abdominal infections.	India  Japan (Drci bKix cūZtbi big Pfizer Limited, India Ges c̄ui emiR K big D <sup>3</sup> GdGmim-tZ উল্লেখ নাই।	cūqRb bvB neaq Arte`b bigÄjy Kiv h̄BtZ ci i	cūqRb bvB neaq Arte`b bigÄjy Kiv nBj
		b) <b>Magnex 2 gm Vial</b>  Sulbactam Sodium USP eq. to Sulbactam 1000 mg + Cefoperazone Sodium IP eq. to Cefoperazone 1000 mg/Vial	-do-	India	RvZiq Jl̄a b̄wZ 2005 tgvZiteK Db̄Z 7U t̄t ki tKibUj GdGmim `wLj Kii tZ ci i bvB neaq Arte`b bigÄjy Kiv h̄BtZ ci i	RvZiq Jl̄a b̄wZ 2005 tgvZiteK Db̄Z 7U t̄t ki tKibUj GdGmim `wLj Kii tZ ci i bvB neaq Arte`b bigÄjy Kiv nBj
13.	Medentech Ltd., Ireland  (Janata Traders)	a) <b>Aquatabs</b>  Sodium Dichlorosocyanurate 33 mg	It is used for water purification	Ireland	Abfgv`b Kiv h̄BtZ ci i	Abfgv`b Kiv nBj
14.	Laboratories CRINEX , France  (Maisha Healthcare)	a) <b>Uvesterol Vitamin ADEC, oral Solution</b>  Synthetic Vitamin A concentrate (Oily form) 300,000 IU (eq. to 176.47 mg) + Ergocalciferol (Vitamin D2) 100,000 IU (eq. to 2.5 mg) + Alpha-tocopherol Acetate (Vitamin E) 0.50 gm + Ascorbic Acid (Vitamin C) 5 gm/100 ml	Indicated for newborn babies (premature babies in particular) and for infants with risk of deficiency or malabsorption of fat soluble vitamins A, D and E and Vitamin C.	France	cūqRb bvB neaq Arte`b bigÄjy Kiv h̄BtZ ci i	cūqRb bvB neaq Arte`b bigÄjy Kiv nBj
		b) <b>Uvedose 100,000 IU Oral Solution in ampoule</b>  Cholecalciferol (Vitamin D3) 2.5 mg (eq. to 100,000 IU) /2 ml Ampoule	Prophylaxis and treatment of vitamin D deficiency	France	Abfgv`b Kiv h̄BtZ ci i	Abfgv`b Kiv nBj
		c) <b>Uvesterol D 1500 IU/ml Oral Solution, 20 ml Bottle with oral syringe.</b>  Ergocalciferol (Vitamin D2) 0.00375 gm (eq. to 150,000 IU)/100 ml	Prevention and treatment of Vitamin D Deficiency in babies and children up to five years.	France	cūqRb bvB neaq Arte`b bigÄjy Kiv h̄BtZ ci i	cūqRb bvB neaq Arte`b bigÄjy Kiv nBj
15.	Swiss Co Services AG, Switzerland  (City Overseas Ltd.)	a) <b>Circadin 2 mg Prolonged release Tablet</b>  Melatonin INN 2 mg	Treatment of primary insomnia characterized by poor quality of sleep in patients who are aged 55 or over.	EMEA	cūqRb bvB neaq m̄m-Gi 238 Zg mfvi Arte`b bigÄjy Kii evi m̄xvš- emij iVLv h̄BtZ ci i	cūqRb bvB neaq Arte`b bigÄjy Kiv nBj

<b>bs</b>	<b>cőZKvitKi big</b>	<b>Jl t̄ai big I tRtbwi K big</b>	<b>ibt`Rbv</b>	<b>CPP/FSC</b>	<b>tUKibKij mve-KugijIi 56 Zg mfvi wmxvš/</b>	<b>mfvi mxvš-</b>
16.	Lilly France, France (International Agencies (Bd) Ltd)	a) <b>Forsteo solution for injection, in 2.4 ml pre-filled pen</b>  Teriparatide,rhPTH(1-34) <sup>1</sup> 20 mcg/80 microliter (250 mcg/ml)	Indicated for the treatment of osteoporosis.	EMEA	<i>Abfgv`b Kiv hBtZ cñti /</i>	<i>Abfgv`b Kiv nBj /</i>
17.	CSL Behring AG, Switzerland (International Agencies (Bd) Ltd.)	a) <b>Rhophylac 300 Solution for Injection (prefilled syringe)</b>  Human Anti-D (RHO) Immunoglobulin 300 mcg/2 ml Prefilled syringe  Human plasma protein ≤ 60 mg + Human anti-D immunoglobulin corresponding to ≥1350 IU ≥270 mcg/2ml Prefilled syringe	Prevention of Rh(D) immunization in Rh(D)- negative women <ul style="list-style-type: none"><li>• Pregnancy/delivery of a Rh(D)-positive baby</li><li>• Abortion/threatened abortion, ectopic pregnancy or hydatidiform mole</li><li>• Transplacental haemorrhage resulting from antenatal haemorrhage, amniocentesis, chorionic biopsy or obstetric manipulative procedures e.g. external version, or abdominal trauma.</li></ul> Treatment of Rh(D)- negative persons after incompatible transfusions of Rh(D)-positive blood or other products containing red blood cells	Switzerland	<i>Abfgv`b Kiv hBtZ cñti /</i>	<i>Abfgv`b Kiv nBj /</i>
18.	S.C Sindan Pharma S.R.L., Romania (Bots Pvt. Ltd.)	a) <b>Eudarym Solution 50 mg/vial for Injection (25mg/ml) (Lyophilisate for solution for injection)</b>  Fludarabine Phosphate 50 mg/Vial (25 mg/ml)	Anti-cancer	Romania Germany	<i>Abfgv`b Kiv hBtZ cñti /</i>	<i>Abfgv`b Kiv nBj /</i>

<b>bs</b>	<b>cūZKvitKi bvg</b>	<b>Jl t̄ai bvg I tRt̄bwii K bvg</b>	<b>ibt̄ Rbv</b>	<b>CPP/FSC</b>	<b>tUKibKij mve-KugijIi 56 Zg mfvi wmxvši</b>	<b>mfvi wmxvši</b>
19.	GalxoSmithKline Biological s.a., Belgium  (GalxoSmithKline Bangladesh Ltd.)	a) <b>Synflorix Suspension for Injection Vaccine</b>  <b>Each dose (0.5 ml) contains :</b> Pneumococcal polysaccharide serotype 1 <sup>1,2</sup> INN 1 mcg Pneumococcal polysaccharide serotype 4 <sup>1,2</sup> INN 3 mcg Pneumococcal polysaccharide serotype 5 <sup>1,2</sup> INN 1 mcg Pneumococcal polysaccharide serotype 6B <sup>1,2</sup> INN 1 mcg Pneumococcal polysaccharide serotype 7F <sup>1,2</sup> INN 1 mcg Pneumococcal polysaccharide serotype 9V <sup>1,2</sup> INN 1 mcg Pneumococcal polysaccharide serotype 14 <sup>1,2</sup> INN 1 mcg Pneumococcal polysaccharide serotype 18C <sup>1,3</sup> INN 3 mcg Pneumococcal polysaccharide serotype 19F <sup>1,4</sup> INN 3 mcg Pneumococcal polysaccharide serotype 23F <sup>1,2</sup> INN 1 mcg <sup>1</sup> adsorbed on aluminium phosphate 0.5 mg Al <sup>3+</sup> <sup>2</sup> Conjugated to protein D (derived from non-typeable Haemophilus influenzae) carrier protein 9-16 mcg <sup>3</sup> Conjugated to tetanus toxoid carrier protein 5-10 mcg <sup>4</sup> Conjugated to diphtheria toxoid carrier protein 3-6 mcg	Active immunization of infants and children from 6 weeks up to 2 years of age against disease caused by Streptococcus pneumoniae serotype 1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F (including sepsis, meningitis, pneumonia, bacteraemia and acute otitis media) and against acute otitis media caused by Non-typeable Haemophilus influenzae.	EMEA	<i>Abfgi`b Kiv hBtZ c̄ti  </i>	<i>Abfgi`b Kiv nBj  </i>
20.	PT. BIO FARMA, Indonesia  (Tajarat Health Care Marketing)	a) <b>BIO-TT Vaccine Injection Ampoule</b>  Purified tetanus toxoid USP 10 Lf/ 0.5 ml	Active immunization against tetanus	Indonesia	<i>RiZiq Jla bñZ 2005 tḡZt̄ek Dbz 7ñU t̄t̄ki tKvbiUi GdGmñ `ñLj Kii‡Z c̄ti bvB ieaq Avte`b bvgÄj Kiv hBtZ c̄ti  </i>	<i>RiZiq Jla bñZ 2005 tḡZt̄ek Dbz 7ñU t̄t̄ki tKvbiUi GdGmñ `ñLj Kii‡Z c̄ti bvB ieaq Avte`b bvgÄj Kiv nBj  </i>
21.	Cadila Healthcare Ltd. India  (Popular Pharmaceuticals Ltd.)	a) <b>Lyssavac N Vaccine</b>  Rabies Inactivated Virus 10 <sup>7.0</sup> LD <sub>50</sub> (as Purified Duck Embryo (PDE) Rabies Vaccine) Equivalent to 2.5 IU per vial (Lyophilized)	Active immunization against rabies	India	<i>RiZiq Jla bñZ 2005 tḡZt̄ek Dbz 7ñU t̄t̄ki tKvbiUi GdGmñ `ñLj Kii‡Z c̄ti bvB ieaq Avte`b bvgÄj Kiv hBtZ c̄ti  </i>	<i>RiZiq Jla bñZ 2005 tḡZt̄ek Dbz 7ñU t̄t̄ki tKvbiUi GdGmñ `ñLj Kii‡Z c̄ti bvB ieaq Avte`b bvgÄj Kiv nBj  </i>

**2.5 Jla mbaSY KigilUi 238 Zg mfvi m̄xv̄S- tḡZt̄eK mbtḡew̄Z c̄. ij i ch̄B Z̄ I KMR-c̄W mn c̄lyiq Arte`b `mLj KwiqtQ t**

bs	c̄ZKvīKi big	Jltai big I tRtbwK big	mb̄Rbv	CPP/FSC	#UKibKij mve-KigilUi 56 Zg mfvi m̄xv̄S/	mfvi m̄xv̄S-
01.	Novo Nordisk A/S, Denmark  (Transcom Distribution)	a) <b>Novo Seven 5 mg (250 KIU) Powder and Solvent for Solution Injection</b>  Eptacog Alfa (Activated) 5 mg/Vial corresponds to 1250 KIU/VIAL, 1mg/ml after reconstitution, 1KIU equals to 1000 IU	Treatment of bleeding episodes and for the prevention of bleeding in those undergoing surgery.	Denmark EMEA	Abfḡ`b Kiv hBtZ cv̄i /	Abfḡ`b Kiv nBj /
		b) <b>Novo Seven 2 mg (100 KIU) Powder and Solvent for Solution for Injection</b>  Eptacog Alfa (Activated) 2 mg/Vial corresponds to 100 KIU/VIAL, 1mg/ml after reconstitution, 1KIU equals to 1000 IU	Treatment of bleeding episodes and for the prevention of bleeding in those undergoing surgery.	Denmark EMEA	Abfḡ`b Kiv hBtZ cv̄i /	Abfḡ`b Kiv nBj /
		c) <b>Novo Seven 1 mg (50 KIU) Powder and solvent for Solution for injection.</b>  Eptacog Alfa (Activated) 1 mg/Vial corresponds to 50 KIU/VIAL, 1mg/ml after reconstitution, 1KIU equals to 1000 IU	Treatment of bleeding episodes and for the prevention of bleeding in those undergoing surgery.	Denmark EMEA	Abfḡ`b Kiv hBtZ cv̄i /	Abfḡ`b Kiv nBj /
02.	OM Pharma, Switzerland  (M/s. Swadesh, Dhaka)	a) <b>Bronchovaxom Adults 40 mg Capsule</b>  Standardized OM-85 lyophilizate corresponding to : Lyophilized bacterial Lysates of : Haemophilus influenzae Diplococcus Pneumoniae Klebsiella Pneumoniae and ozaenae Staphylococcus aurcus Streptococcus pyogenes and viridans Neisseria catarrhalis eq to 7mg.	Immunotherapy, prevention of recurrent chronic bronchitis, Tonicitis, Pharyngitis and bacterial resistance to conventional therapy.	Switzerland	c̄qRb b̄B neaq Arte`b bigAj̄ Kiv hBtZ cv̄i /	c̄qRb b̄B neaq Arte`b bigAj̄ Kiv nBj /
03.	Berna Biotech Korea Corp.,Korea  (Renata Ltd.)	a) <b>Quinvaxem Vaccine</b>  DTP-HePB-Hib Fully Liquid combined Vaccine	Prevent diphtheria, tetanus, pertussis, hepatitis B and haemophilus influenzae type B.	Korea	RiZiq Jla b̄W 2005 tḡZt̄eK Db̄Z 7JU t̄tki tKiblUi GdGmim `mLj KwīZ cv̄i b̄B neaq Arte`b bigAj̄ Kiv hBtZ cv̄i /	RiZiq Jla b̄W 2005 tḡZt̄eK Db̄Z 7JU t̄tki tKiblUi GdGmim `mLj KwīZ cv̄i b̄B neaq Arte`b bigAj̄ Kiv nBj /

2.6 *AbtqeevY<sup>2</sup> c`W JIa ibqSY KigUj 238 Zg mfvi m×vš- tgZteK tUKibK<sup>v</sup>j mve-KigUj mfvq gZig<sup>v</sup>Zi Rb<sup>v</sup> Dc<sup>v</sup>cb Kiv nq t*

<i>bs</i>	<i>cōZKvitKi bvq</i>	<i>JItaibv I tRtbvK bvq</i>	<i>Abt<sup>v</sup>Rbv</i>	<i>CPP/FSC</i>	<i>m×vš-Gi 238 Zg mfvi m×vš-</i>	<i>tUKibK<sup>v</sup>j mve-KigUj 56 Zg mfvi m×vš</i>	<i>mfvi m×vš-</i>
1.	M/s. Schering Oy Turku, Finland  (M/s. Rubeus & Rasa Pvt. Ltd.)	a) <b>Jadelle 75mg Implant</b>  Levonorgestrel USP 75mg/Implant	Contraception	Finland Spain France	D <sup>3</sup> JIaUj we <sup>v</sup> q cwi evi cwi Kí bv Awa`B <sup>v</sup> i i <sup>v</sup> iK bs- ccA/mwmGmWic/5364, Zwil t 14-10-2009 Gi c <sup>v</sup> i i cwi tc <sup>v</sup> tz mfvq Avtj vPbv nq/ <b>Jadelle 2 i W neekó, eZg<sup>v</sup>tb</b> t <sup>v</sup> tk <b>Jadelle Gi GKU Utqj</b> cwi Pvj Z nB <sup>v</sup> ZtQ GLb ch <sup>v</sup> - Utqj Gi dj vdj mtš-vl RbK/ c`W 5 erm <sup>v</sup> i i GKU `xN <sup>©</sup> tgqv`x c <sup>v</sup> W nI qvq ht_ó gj- mvk <sup>v</sup> x/ tUKibUK <sup>v</sup> j mve-KigUj mfvq clyivq Dc <sup>v</sup> cb ce <sup>v</sup> gZigZmn JIa ibqSY KigUj <sup>v</sup> Z Dc <sup>v</sup> cb Kiv eri Rb <sup>v</sup> ej v nBj /	Abtqeev b Kiv hB <sup>v</sup> Z c <sup>v</sup> i /	Abtqeev b Kiv nBj /

## 2.7 Proposed Product for Import (Veterinary)

bs	cōZKvičKi bvg	Jl̄tai bvg I tR̄tbwīK bvg	ibt`Rbv	CPP/FSC	tUKibK'ij mve-KigwJi 56 Zg mfvi m̄xvš-	mfvi m̄xvš-
01.	S.P. Veterinaria S.A., Spain  (Haychem (Bangladesh) Ltd.)	a) <b>Endoex Oral (Vet)</b>  Closantel BP 0.05 gm/ml	Broad spectrum antiparasitic agent used against several species and developmental stages of trematodes, nematodes and arthropods of cattle, sheep, goats, horses and camels.	Spain	cōq̄Rb b̄B m̄araq Avte`b bvgÄy Kiv h̄BtZ c̄ti	cōq̄Rb b̄B m̄araq Avte`b bvgÄy Kiv nBj
02.	Han Dong Co. Ltd., Korea  (Precisa Techno Trade)	a) <b>Amitop-S Injection (Vet) (Amino acids &amp; vitamins)</b>  L-Lysine 170 mg L-Methionine 51 mg L-Threonine 78.2 mg L-Tryptophan 34 mg L-Arginine 142.5 mg L-Leucine 187 mg L-Isoleucine 85 mg L-Valine 136 mg L-Phenylalanine 119 mg Glycine 375 mg Sodium L-Glutamate 375 mg Vitamin B <sub>1</sub> 10 mg Vitamin B <sub>2</sub> 5 mg Vitamin B <sub>6</sub> 10 mg Vitamin B <sub>12</sub> 5 mcg Calcium Pantothenate 10 mg Nicotinamide 300 mg Sodium Acetate Trihydrate 750 mg Calcium Chloride Dihydrate 200 mg Potassium Chloride 200 mg Magnesium Chloride Hexahydrate 200 mg Glucose 5000 mg/100 ml	Hypovitaminosis in all kind of animal, growth troubles, anorexia, asthenia, Convalescence and prophylactic treatment.	Korea	Abtgr`b Kiv h̄BtZ c̄ti	Abtgr`b Kiv nBj
		b) <b>Floshot 300 Injection (Vet)</b>  Florfenicol USP 300 mg/ml	Treatment of BRD, Pneumonia, Pododermatitis, PRDC, Colibacillosis, salmonellosis, glasser's disease etc.	Korea	cōq̄Rb b̄B m̄araq Avte`b bvgÄy Kiv h̄BtZ c̄ti	cōq̄Rb b̄B m̄araq Avte`b bvgÄy Kiv nBj

bs	cōZKvičKi bvg	Jl̄tai bvg I tR̄tbwi K bvg	ibt`Rbv	CPP/FSC	tUKibK'ij mve-KigwJi 56 Zg mfvi m̄xvš	mfvi m̄xvš-
Han Dong Co. Ltd., Korea (Precisa Techno Trade)	c) <b>D-Pron Injection (Vet)</b>  Sulpyrine USP 400 mg/ml	MMA, Pneumonia, Mastitis, Metritis, Agalactia and other inflammatory disease.	Korea	Abtgr`b Kiv h̄BtZ c̄ti	Abtgr`b Kiv nBj	
	d) <b>Floshot Oral Solution (Vet)</b>  Florfenicol USP 100 gm/litre	Treatment of bacterial disease caused by sensitive to florfenicol like as mycoplasma spp. E.coli, Salmonella, Streptococcus, Pasteurella, Staphylococcus spp.	Korea	c̄qRb b̄B m̄ainq Avte`b bigÄj Kiv h̄BtZ c̄ti	c̄qRb b̄B m̄ainq Avte`b bigÄj Kiv nBj	
03.	Investigaciones Quimicas Y Farmaceuticals SA, Spain (Vet Healthcare Agency)	a) <b>Desinkap Oral Liquid (Vet)</b>  Glyoxol 40% 17 g + Formaldehyde 40% 15 g + Benzalkonium Chloride 80% 13 g + Glutaraldehyde 50% 50 g/liter	Surface Disinfectant for vehicles, hatching rooms, livestock stables, dairy rooms.	Spain	Abtgr`b Kiv h̄BtZ c̄ti	Abtgr`b Kiv nBj
04.	Izo S.P.A. Via A Bianchi-9, Italy (Jayson Agrovet Ltd.)	a) <b>IZOVAC MAREK BIVALENT (HVT+RISPENS) (Vet)</b>  Cell associated live frozen herpes turkey virus of Marek's Disease Strain FC 126 (serotype 3) 2000 PFU + Cell associated live frozen Marek's Disease Virus strain Rispense CVI988 (serotype 1) 2000 PFU/0.2 ml dose	Cell associated frozen live vaccine against marek's disease.	Italy	Abtgr`b Kiv h̄BtZ c̄ti	Abtgr`b Kiv nBj
05.	Bomac Laboratories Ltd., Newzealand (Remax Corporation Ltd.)	a) <b>Terminator Liquid (Vet)</b>  Glutaraldehyde 50%, 150 gm + Synthecol CC 50 (Cocobenzylidemethylammonium Chloride) (Quaternary Ammonium Chloride) 100 gm/Litre	For all surface disinfection & aerial fogging of livestock premises especially for hygiene programmes associated with calfrearing and pig and poultry farms.	Newzealand Australia	Abtgr`b Kiv h̄BtZ c̄ti	Abtgr`b Kiv nBj
06.	Kela Laboratoria NV, Belgium (Ceder Bangladesh Ltd.)	a) <b>Amprolium 20% Water Soluble Powder (Vet)</b>  Amprolium Hydrochloride 200 mg/Kg	Treatment and control of coccidiosis.	Belgium	c̄qRb b̄B m̄ainq Avte`b bigÄj Kiv h̄BtZ c̄ti	c̄qRb b̄B m̄ainq Avte`b bigÄj Kiv nBj
07.	DEC International NZ Ltd., Newzealand (Janata Traders)	a) <b>CIDR Cattle Insert (Vet) (Intra vaginal progesterone insert)</b>  Progesterone USP 1.38 gm/Insert	To advance the date of first estrus in postpartum beef cows and in peripubertal beef heifers. To synchronize estrus in beef cows, and in beef and dairy heifers.	Newzealand	Abtgr`b Kiv h̄BtZ c̄ti	Abtgr`b Kiv nBj
08.	EWABO Chemikalien GmbH & Co. KG, Germany (Univet Limited)	a) <b>Mentofin (Vet) Oral liquid</b>  Eucalyptus oil + Menthol 200 gm/kg	To increase mucous secretion in case of problem with respiratory system, reduce post vaccination reaction, to cool the dry mucous membranes & to reduce inflammation by inhibiting release of inflammatory mediators.	Germany	c̄qRb b̄B m̄ainq Avte`b bigÄj Kiv h̄BtZ c̄ti	c̄qRb b̄B m̄ainq Avte`b bigÄj Kiv nBj

bs	cōZKvičKi bvg	Jl̄tai bvg I tRtbwīK bvg	ibt`Rbv	CPP/FSC	tUKibK'ij mve-KigwJi 56 Zg mfvi m̄xvš-	mfvi m̄xvš-
09.	Shanghai Veterinary Pharmaceuticals, China  (Advance Animal Science Co. Ltd.)	a) <b>Auflosyn Injectable Solution</b>  Ofloxacin 100mg/ml	Broad spectrum antibacterial and is used mainly for bacterial as well as microbacterial infection in animals.	China	cōqRb bvB neaq Avte`b bvgÄj Kiv hBtZ cti i	cōqRb bvB neaq Avte`b bvgÄj Kiv nBj
		b) <b>Nikang Injection</b>  Florfenicol 300mg/ml	Broad spectrum synthetic antibiotic active against most gram positive and gram negative bacteria. It can be used to control bacterial disease of swine, chicken and fish caused by susceptible bacteria.	China	cōqRb bvB neaq Avte`b bvgÄj Kiv hBtZ cti i	cōqRb bvB neaq Avte`b bvgÄj Kiv nBj
		c) <b>Amisyn Injection</b>  Amikacin 100 mg/ml	Bactericidal action against Escherichia, brucella, calymmatobacterium, citrobacter, enterobacter, francisella, klebsiella, proteus, etc.	China	cōqRb bvB neaq Avte`b bvgÄj Kiv hBtZ cti i	cōqRb bvB neaq Avte`b bvgÄj Kiv nBj
		d) <b>Lomesyn Injection</b>  Lomefloxacin 100 mg/ml	Antibacterial and used mainly for bacterial infections in animals. It is especially indicated for use in treating respiratory disease, gastrointestinal and urinary tract infections.	China	cōqRb bvB neaq Avte`b bvgÄj Kiv hBtZ cti i	cōqRb bvB neaq Avte`b bvgÄj Kiv nBj
10	Adisseo, France  (Advance Animal Science Co. Ltd.)	a) <b>Rovabio Excel AP T-Flex Powder</b>  Endo-1, 4-β-xylanase : No EC 3.2.1.8 Endo-1, 3(4)- β -glucanase: No EC 3.2.1.6 + 17 other enzyme activities, including cellulose (endo-1,4- β -glucanase)	Improves the nutritional value of feeds containing different types of cereals (wheat, maize, barley, triticale, rye, oats) and oilseed meals, reduces gut viscosity, reduces the free ammonia concentration in the litter.	France	Abfgv`b Kiv hBtZ cti i	Abfgv`b Kiv nBj
		b) <b>Rovabio Max AP Powder</b>  Endo-1, 4-β-xylanase : No EC 3.2.1.8 Endo-1, 3(4)- β -glucanase: No EC 3.2.1.6 6 Phytase No EC 3.1.3.26 Other enzyme activities : cellulases	-do-	France	Abfgv`b Kiv hBtZ cti i	Abfgv`b Kiv nBj
		c) <b>Rovabio Excel LC Liquid</b>  Endo-1, 4-β-xylanase ≥ 5500 U Visco / g ..... 3.00% eq. to 350 Units AXC/g Endo-1, 3(4)- β -glucanase ≥ 500 AGL U/g ..... 3.00%	-do-	France	cōqRb bvB neaq Avte`b bvgÄj Kiv hBtZ cti i	cōqRb bvB neaq Avte`b bvgÄj Kiv nBj

bs	cōZKvičKi bvg	Jl̄tai bvg I tRtbwīK bvg	ibt`Rbv	CPP/FSC	tUKibK'ij mve-KigwJi 56 Zg mfvi wmxvš	mfvi wmxvš-
11.	Merial Select Inc., USA  (Advance Animal Science Co. Ltd.)	a) <b>VAXXITEK HVT + IBD Vaccine</b>  Each dose of vaccine contains: Marek's Disease vectored IBD recombinant virus, at least 3,880 PFU	This vaccine is recommended for vaccination of 18 to 19 days old embryonated chicken eggs. For in Ovo vaccination, only healthy embryonated eggs should be used.	USA	Abtgv`b Kiv hBtZ c̄ti	Abtgv`b Kiv nBj
12.	Kemira ChemSolutions bv, The Netherlands  (Advance Animal Science Co. Ltd.)	a) <b>Kemira Mould Control SP1 Granulate</b>  Calcium Propionate (on dried matter) min. 985gm/Kg	Kemira Mould control SP1 prevents growth of mould and other harmful microbes in different types of feedstuffs. It also prevents recontamination after production of the compound feed	The Netherlands	Abtgv`b Kiv hBtZ c̄ti	Abtgv`b Kiv nBj
		b) <b>Kemira Mould Control LP3 NC Liquid</b>  Propionic Acid 242gm/Kg Ammonium Propionate 563gm/Kg Benzoic Acid 10gm/Kg Surfactant 2 gm/Kg	Kemira Mould control LP3 NC has a strong anti mould effect in compound feeds and raw materials and prevents recontamination of mould after product of the feed	The Netherlands	Abtgv`b Kiv hBtZ c̄ti	Abtgv`b Kiv nBj
		c) <b>Kemira Rumen Pro SP1 Powder/Granulate</b>  Propionic Acid min. 662 gm/Kg	Kemira Rumen Pro SP1 has been developed to increase rumen propionate content without lowering rumen pH	The Netherlands	c̄qRb bB w̄aq Avte`b bvgÄj Kiv hBtZ c̄ti	c̄qRb bB w̄aq Avte`b bvgÄj Kiv nBj
		d) <b>Kemira Pro GIT SF1 Powder</b>  Formic Acid min. 306 gm/Kg Acetic Acid min. 123 gm/Kg Lactic Acid min. 101 gm/Kg Citric Acid min. 72 gm/Kg	The organic acids in Kemira Pro GIT SF1 reduce crop pH and growth of pathogenic microbes in the gastrointestinal (GI tract). Reduced pathogenic microbial growth in the GI tract also severs nutrients for the animal.	The Netherlands	Abtgv`b Kiv hBtZ c̄ti	Abtgv`b Kiv hBtZ c̄ti
		e) <b>Kemira Bacteria Control SF1 Powder</b>  Formic Acid min. 408 gm/Kg Propionic Acid Min 103 gm/Kg Citric Acid min. 130 gm/Kg	Dry Antibacterial agent	The Netherlands	c̄qRb bB w̄aq Avte`b bvgÄj Kiv hBtZ c̄ti	c̄qRb bB w̄aq Avte`b bvgÄj Kiv nBj
13.	Kemira ChemSolution bv, The Netherlands  (Advance Animal Science Co.Ltd.)	a) <b>Kemira Mould Control SP1</b>  Calcium Propionate INN			c̄qRbq Z_w̄ br_vKvq Avte`b bvgÄj Kiv hBtZ c̄ti	c̄qRbq Z_w̄ br_vKvq Avte`b bvgÄj Kiv nBj

bs	cōZKvičKi bvg	Jl̄tai bvg I tRtbwīK bvg	ibt`Rbv	CPP/FSC	tUKibK'ij mve-KigwJi 56 Zg mfvi wmxvš	mfvi wmxvš-
14.	DaOne Chemical Co. Ltd., Korea (Wilts Marketing Ltd.)	a) <b>DaOne E+ Selenium Oral Liquid</b>  Vitamin E (Alpha tocopherol acetate) 17 gm + Selenium (Sodium Selenite) 2 gm/Litre	Improvement of feed efficiency and promotion of growth rate.	Korea	Abtgr`b Kiv hBtZ cti /	Abtgr`b Kiv nBj /
		b) <b>DaOne ADEK Oral Liquid</b>  Vitamin A 10,000,000 IU + Vitamin D <sub>3</sub> 200000IU + Vitamin E 4000 IU + Vitamin K <sub>3</sub> 132 mg/Litre	Prevention of stress and vitamin A, D, E, K deficiency,	Korea	Abtgr`b Kiv hBtZ cti /	Abtgr`b Kiv nBj /
		c) <b>DaOne Tilmiro Solution</b>  Tilmicosin Phosphate 100 gm/litre + Xylo-oligosaccharide qs.	Treatment of respiratory diseases caused by sensitive bacterial to tilmicosin as follows: Pig – Pasteurella multocida, Actinobacillus pleuropneumiae, Mycoplasma hyopneumoniae Poultry : Mycoplasma galisepticum, Mycoplasma synoviae.	Korea	cōqirb bvb weaq Avte`b bigÄj Kiv hBtZ cti /	cōqirb bvb weaq Avte`b bigÄj Kiv nBj /
15.	Dong Bang Co. Ltd., Korea (Wilts Marketing Ltd.)	a) <b>Control Injection</b>  Fertireline Acetate 20 mcg + Sodium Chloride 3.6 mg/ml	Treatment of ovarian follicular cysts, delayed ovulation, anovulation.	Korea	cōqirb bvb weaq Avte`b bigÄj Kiv hBtZ cti /	cōqirb bvb weaq Avte`b bigÄj Kiv nBj /
		b) <b>Gonadon Injection</b>  Gonadoreline (as acetate) BP 100 mcg + Benzyl Alcohol BP 10 mg + Sodium Chloride BP 7.47 mg/ml	For the treatment of ovarian follicular cysts, delayed ovulation, anovulation. For the prevention and treatment of abnormal oestrus, synchronization of oestrus.	Korea	Abtgr`b Kiv hBtZ cti /	Abtgr`b Kiv nBj /
		c) <b>Dinorin Injection</b>  Cloprostenol (as sodium) BP 100 mcg/ml	Cattle: Control the time of oestrus and ovulation. Synchronization of the oestrus cycle. Treat cattle which have a functional corpus luteum but do not express behavioral oestrus. Termination of abnormal pregnancy. Treatment of chronic metritis and pyometra. Routine use in the early postpartum period. Luteal cysts. Pigs: Induction of parturition.	Korea	Abtgr`b Kiv hBtZ cti /	Abtgr`b Kiv nBj /
		d) <b>Fortis Injection</b>  Flunixin Meglumine 50 mg/ml	Horse : For the alleviation of inflammation and pain associated with musculoskeletal disorder Cattle : For the control of pyrexia associated with bovine respiratory disease and endotoxemia and acute bovine mastitis. For the control of inflammation in endotoxemia Swine : Mastitis, metritis, agalactia syndrome, and acute inflammatory conditions and pyrexia due to pneumonia or diarrhea.	Korea	Abtgr`b Kiv hBtZ cti /	Abtgr`b Kiv nBj /

<b>bs</b>	<b>cōZKvičKi bvg</b>	<b>JItai bvg I tRtbwi K bvg</b>	<b>ibt`Rbv</b>	<b>CPP/FSC</b>	<b>tUKibK'ij mve-KugUi 56 Zg mfvi m×vši</b>	<b>mfvi m×vš-</b>
16.	Laboratories Ovejero SA, Spain (Tajarat Animal Care)	a) <b>Inmugal V.P. Hitchner B-1</b>  Active virus of Newcastle disease, attenuated strain Hitchner B1 10 <sup>6</sup> EID <sub>50</sub> /Dose	Active immunization of chickens and adult hens against Newcastle disease.	Spain	cōqRb bvB neaq m×vš-Gi 238 Zg mfviq Avte`b bvgÄý Kii evi m×vš- enyj iLv hBtZ cti	cōqRb bvB neaq Avte`b bvgÄý Kiv nBj
17.	KBNP Inc, Korea (Tajarat Animal Care)	a) <b>Himmvac ND Oil Vaccine</b>  Inactivated ND virus (B1) -15% Marcol 52-58% Montanide-7%	Prevention Newcastle disease in poultry.	Korea	cōqRb bvB neaq m×vš-Gi 238 Zg mfviq Avte`b bvgÄý Kii evi m×vš- enyj iLv hBtZ cti	tUKibK'ij mve-KugUi cieZmfvq ctiq Dc`vcibi Rb` ej v nBj
	b) <b>Himmvac IB ND Combined Live Vaccine</b>  Newcastle Disease Virus (B1) infected fluid-30% IB virus (H120) infected fluid – 15% Stabilizer (LPG) -20% Penicillin – 200IU/ml Streptomycin – 200mcg/ml	Active immunization against Infectious Bronchitis and Newcastle Disease.		Korea	cōqRb bvB neaq m×vš-Gi 238 Zg mfviq Avte`b bvgÄý Kii evi m×vš- enyj iLv hBtZ cti	tUKibK'ij mve-KugUi cieZ <sup>®</sup> mfviq ctiq Dc`vcibi Rb` ej v nBj
	c) <b>Himmvac ND High Vaccine</b>  Amount per dose Newcastle Disease Virus (Ulster) infected fluid – 10 <sup>6.0</sup> EID <sub>50</sub> /dose Stabilizer (LPGG) – 20% Penicillin – 200IU/ml Streptomycin – 200mcg/ml	Prevention of Newcastle disease in chickens.		Korea	Abfgv`b Kiv hBtZ cti	Abfgv`b Kiv nBj
	d) <b>Himmvac IB-ND Combined oil vaccine</b>  Inactivated IB virus (H-120) -20% Inactivated ND virus (B1) – 15% Marcol 52 – 58% Montanide – 7%	Prevention Infectious Bronchitis Newcastle disease.		Korea	cōqRb bvB neaq m×vš-Gi 238 Zg mfviq Avte`b bvgÄý Kii evi m×vš- enyj iLv hBtZ cti	tUKibK'ij mve-KugUi cieZ <sup>®</sup> mfviq ctiq Dc`vcibi Rb` ej v nBj
	e) <b>Himmvac IB-ND-EDS Combined oil Vaccine</b>  Inactivated IB virus (H-120) – 20% Inactivated ND virus (B1) – 15% Inactivated EDS'76 virus – 15%	Prevention of Infectious Bronchitis Disease & Egg Drop Syndrome 76 in poultry.		Korea	Abfgv`b Kiv hBtZ cti	Abfgv`b Kiv nBj

bs	cōZKvičKi bvg	Jl̄tai bvg I tRtbwīK bvg	ibt`Rbv	CPP/FSC	tUKibK'ij mve-KigwJi 56 Zg mfvi m̄xvš-	mfvi m̄xvš-	
	KBNP Inc, Korea (Tajarat Animal Care)	<p>f) <b>Himmvac Newcastle Live Vaccine (B1 Strain)</b>            Amount per 1000 doses            Newcastle Disease Virus (B1) infected fluid – 50%            Stabilizer (LPGG) – 50%            Penicillin – 200IU/ml            Kanamycin – 100mcg/ml            Streptomycin – 200mcg/ml</p> <p>g) <b>Himmvac Fowl Pox Vaccine</b>            Fowl Pox virus infection Fluid 30%            Stabilizer 70%            Penicillin 200 IU/ml            Kanamycin 100 mcg/ml            Streptomycin 200 mcg/ml</p> <p>h) <b>Himmvac ILT Live vaccine</b>            Infectious Laryngotracheitis virus infected fluid            50%            Stabilizer 50%            Streptomycin 200 mcg/ml            Penicillin 200 IU/ml</p>	Prevention of Newcastle Disease in Chickens.  For vaccination of healthy chickens against fowl pox.  For vaccination of healthy chicken against infections laryngotracheitis.	Korea  Korea  Korea	<i>cōqRb bvb neaq m̄mm-Gi 238</i> <i>Zg mfvi Arte`b bvgÄy Kiv eri</i> <i>m̄xvš- enj ivlv hibtz c̄ti /</i>	<i>cōqRb bvb neaq Arte`b</i> <i>bvgÄy Kiv nBj /</i>	<i>Abfgv`b Kiv hibtz c̄ti /</i>
18.	Laboratories Hipra SA, Spain (Nasco Agro Product)	<p>a) <b>Hipra Gumboro-GM97 Live Freeze dried Vaccine</b>            Live Infectious Bursal Disease Virus, strain GM97  <math>\geq 10^2 \text{ EID}_{50}</math></p>	To prevent Gumboro Disease.	Spain	<i>Abfgv`b Kiv hibtz c̄ti /</i>	<i>Abfgv`b Kiv nBj /</i>	

<i>bs</i>	<i>cōZKviłKi bıg</i>	<i>Jl̄tai bıg I tR̄tbwı K bıg</i>	<i>ıbt̄ Rbv</i>	<i>CPP/FSC</i>	<i>tUKibK'ıj mve-KıgıJı 56 Zg mfvi m̄xvısl̄</i>	<i>mfvi m̄xvısl̄</i>
19.	Komipharma International Co. Ltd., Korea  (Rafique Medicine)	<p>a) <b>Amcicoli Powder</b>  Ampicillin Trihydrate 200 gm + Colistin Sulfate 500,000,000 IU/Kg</p> <p>b) <b>Komicelase Powder</b>  Celluase 1,000,000 Unit + <math>\alpha</math>-Amylase 40,000 Unit + <math>\beta</math>-Amylase 2,550 Unit + Protease 1,25,000 Unit/Kg</p> <p>c) <b>Diloxa Udder Ointment</b>  Each 1 Syinge (7.5 g) contains : Dicloxacillin Sodium 500 mg</p>	For the prevention and treatment of bacterial disease which are sensitive to ampicillin and colistin. (1) Cattle (under 6 months), Pig : Salmonella, E.coli, pasteurella (2) Poultry : Mycoplasma, Haemophilus, E.coli, salmonella, other streptococcus, staphylococcus.	Korea	<i>Abtgr`b Kiv h̄BtZ cıtı /</i>	<i>Abtgr`b Kiv nBj /</i>
20.	Laboratoire Sogeval, France  (Century Agro Limited)	a) <b>Disinfectant TH4+</b>  <u>Each 1 Litre contain</u> Didecyl dimethyl ammonium chloride 18.75 g Diocetyl dimethyl ammonium chloride 18.75 g Octyl decyldimethyl ammonium chloride 37.50 g Alkyl dimethylbenzyl ammonium chloride 50.00 g Glutardialdehyde 62.50 gm	Disinfectant	France	<i>Abtgr`b Kiv h̄BtZ cıtı /</i>	<i>Abtgr`b Kiv nBj /</i>

**2.8 ।।tgaewZ c`॥ JIa ।।qšY KugUj 238 Zg mfvi ।।m×vš- tgiZiteK tUKibKij mve-KugUj mfviq gZigtZi Rb` cbyiq Dc`lcb Kiv nq t**

bs	cÖZKvitKi bvıg	JIłai bvıg I tRtbıwK bvıg	।।t`Rbv	CPP/FSC	tUKibKij mve-KugUj 56 Zg mfvi ।।m×vš	mfvi ।।m×vš-
1.	Intervet India Pvt. Ltd., India  (Bengal Overseas Ltd.)	a) <b>Decivac FMD DOE Vial Vaccine</b>  Inactivated FMD Virus Serotypes O.A. and Asia 1 at least 3PD <sub>50</sub> per serotype	Immunization of healthy livestock as an aid in the control of FMD.	India	।।m×vš-Gi ।।m×vš tgiZiteK tftUti bvix JIa Avg`vbxi tPitc thvR`Dbz 24nU t`tki tKibUj GdGmim `mlj KwiZ ci i bvB neavq Avte`b bvıgÄy Kiv hvBtZ ci i /	।।m×vš-Gi ।।m×vš tgiZiteK tftUti bvix JIa Avg`vbxi tPitc thvR`Dbz 24nU t`tki tKibUj GdGmim `mlj KwiZ ci i bvB neavq Avte`b bvıgÄy Kiv nBj /

**2.9 Proposed Product for Import (Surgical Equipment & Others)**

Sl. No	Name of the Manufacturer	Name of the Product	Indication	FSC	tUKibKij mve-KugUj 56 Zg mfvi ।।m×vš	mfvi ।।m×vš-
01.	Becton Dickinson SA, Spain  (Mix Medical, 9/A Motaleb Tower, 3 <sup>rd</sup> Floor), 8/2, Paribagh, Hatirpool, Dhaka)	a) <b>BD Soloshot 1 ml AD Syringes 22GX1</b>  Auto Disable Syringe 1 ml	Syringe		cqiqRbq wdktmj mwUldtKU `mlj bvıq Avte`b bvıgÄy Kiv hvBtZ ci i /	cqiqRbq wdktmj mwUldtKU `mlj bvıq Avte`b bvıgÄy Kiv nBj /
02.	Convex Latex Pvt. Ltd., India  (UGI Trade Co., )	a) <b>Natural rubber Latex Condom</b>	Contraception and for prophylactic purpose.	India	cqiqRb bvB neavq Avte`b bvıgÄy Kiv hvBtZ ci i /	cqiqRb bvB neavq Avte`b bvıgÄy Kiv nBj /

<i>bs</i>	<i>meei Y</i>	<i>tUKibKij mve-Kigui</i> 56 Zg mfvi m×vS/	<i>mfvi m×vS-</i>
2.10	<p><i>tgmwmqivk evsj vt' k wj t Gi Avte` Z</i> Made for F. Hoffmann La Roche Ltd. Switzerland by Chugai Pharma Manufacturing Company Ltd. Japan <i>Gi Drcw` Z Actemra 400 mg/20 ml Vial (Tocilizumab 400 mg/20 ml)</i> <i>bigrq c` wji tiRt÷kb Avte` b clytePbri Rb` Avte` b Kwi qvtQ/</i></p> <p>উলেখ্য যে, বিগত ২২-10-2009 <i>Zwi tL AbjøZ Jia wbqSY Kigui</i> 238 Zg mfviq (1) <b>Actemra 80 mg/4ml Vial (Tocilizumab 20 mg/ml) Ges (2) Actemra 200 mg/10 ml Vial (Tocilizumab 20 mg/ml) Abjgr` b Kiv nq Ges <b>Actemra 400 mg/20 ml Vial (Tocilizumab 20 mg/ ml)</b> <i>Gi tiRt÷kb Avte` b bigÄy Kiv nq/</i></b></p> <p>প্রতিষ্ঠানটি তাহাদের আবেদনে উলেখ্য করিয়াছে যে, Actemra <i>Gi</i> Standard dose, 8 mg per Kg body weight, <i>dtj cØB eq` tiMf i Rb` 400 mg/20 ml cØq/Rb / Moderate to severe active rheumatoid arthritis (RA) wPKrmvi Rb` Zimn` i Avte` b wetePbri Rb` Abjiva Kwi qvtQ/</i></p> <p><i>cØzbvbiUi Avte` tbi cwtcØtZ Avte` Z Actemra 400 mg/20 ml Vial (Tocilizumab 20 mg/ ml) Gi tiRt÷kb Avte` b Abjgr` tbi weI tq m×vS- MøtYi Rb` Dc` vcb Kiv nq/</i></p>	<i>Abjgr` b Kiv hvBtZ cifti /</i>	<i>Abjgr` b Kiv nBj /</i>
2.11	<p>M/s. Advance Animal Science Co. Ltd., M/s. Merial Select Inc. <i>cØz÷vbi USA KZR.Drcw` Z c` h_v (1) Galivac AE+ FP (Freeze Dried live Vaccine) (Avian Encephalomyelitis Virus at list ... 2.7 Log 10 EID<sub>50</sub> + Fowl Pox Virus at least 2.2 log 10EID<sub>50</sub>) (2) Trovac AIV H5 Vaccine (Fowl pos vectored ai recombinant virus at least 2.2 log 10 TCID<sub>50</sub>/Dose of freeze dried vaccine), (3) Gallimune Flu H5N9 Vaccine (Inactivated Avian Influenza virus H5N9 strain ≥10HIU) and (4) Gallimune 208 ND + Flu H9 M, E Vaccine (Inactivated Avian Influenza virus H9N2 strain ≥10HIU + Inactivated Newcastle disease virus, Ulster 2C strain ≥10HIU) <i>Avg` vbx tiRt÷k` tbi Avte` b clytePbri Rb` Avte` b Kwi qvtQ/ weMZ 22-10-2009 Zwi tL AbjøZ Jia wbqSY Kigui</i> 238 Zg mfviq D<sup>3</sup> c` mgr cØq/Rb bvB weaq bigÄy Kiv nq/</i></p> <p>প্রতিষ্ঠানটি উলিংচথিত ভ্যাকসিনসমূহ এভিয়ন ইনফ্রয়েজ্বা রোগ প্রতিরোধে নিরাপদ বলিয়া তাহাদের পত্রে উলেখ্য করিয়াছে। এভিয়ন ইনফ্রয়েজ্বা ভ্যাকসিন ব্যবহারের কিছু ইতিবাচক দিক পত্রে উলেখ্য করা হইয়াছে। প্রতিষ্ঠানটির ব 3e "Abjgrq t` tk Gwfqvb BbdqÄvi gnugivx weivR Kivq tcme` wkti i gñZ mñaz nBqvtQ, tcme` wkti K i gñvi Rb` Gwfqvb BbdqÄvi f'vkmib wekli Ri " ix/</p> <p>উলিংচথিত ভ্যাকসিনসমূহের রেজিস্ট্রেশনের আবেদন অনুমোদনের বিষয়ে <i>m×vS- MøtYi Rb` Dc` vcb Kiv nq/</i></p>	<i>cØq/Rb bvB weaq Avte` b bigÄy Kiv hvBtZ cifti /</i>	<i>cØq/Rb bvB weaq Avte` b bigÄy Kiv nBj /</i>

<i>bs</i>	<i>meiY</i>	<i>#UKibKij mve-KugUj 56 Zg mfvi m×vš</i>	<i>mfvi m×vš-</i>
2.12/	<p><i>tgmwm®ZRviZ tnj &amp; tKqvi gvtKls Drug Control Committee (DCC) Gi 235 Zg mfiq (10 Gicj, 2007 Zwi tL AbjøZ) Anti-Cancer Drug রেজিস্ট্রেশন সংক্রান্ত- m×vš- ev~evgtbi Rb~ Avte`b Kii qitQ/ cñZomU Zmvt`i ctí উল্লেখ করিয়াছে যে, বিশ্বের বিভিন্ন দেশে নতুন নতুন ক্যাপ্সার চিকিৎসার ওষধ তৈরী হইতেছে কিন্তু আমদানী জটিলতার কারণে বাংলাদেশের ক্যাপ্সার আক্রান্ত- tivMiv GB mKj JIa t_tK evtA nBtZtQ/ cñZomU JIa ibqSY KugU (Wmim) Gi 235 Zg mfiq Anti-Cancer Drug mnRj f' Kii evi Rb~ Zwi Krf3 24 ll t`k nBtZ Avg`vbi `eaZvi nel qiu KvHkix Kii evi Rb~ Abjiva Kii qitQb/</i></p> <p><b>JIa ibqSY KugU (Wmim) Gi 235 Zg mfiq Anti-Cancer Drug Gi tIRt-kb nel tq Avtj vPbv / m×vš- t</b></p> <p><i>eijs vt`k digmDiUKij Bt~vUfm®Gtmwmtqkb evRiti `jvtiM KivYvi Pukrmvi JIamgtni mnR I mijf ctc~Zv ibioZ Kii evi j tpt`ibtgié tcñlvcU eYBv ceR Avte`b tck Kti   Avte`b Abjyt i KivYvi Pukrmvi JIamgtni ctk®Zr®`kmn weifbae t`k nBtZ A%ধ মাধ্যমে ইতঃপূর্বে বাংলাদেশের বাজারে প্রবেশ করিলেও বর্তমানে এই সুযোগ নাই। ফলে ক্যাপ্সার রোগে আক্রান্ত-MtYi Pukrmvi AebiZ NuñZtQ ejq qu Avte`b `ver Kiv nq/</i>  <i>এই ক্ষেত্রে উল্লেখ্য যে, ক্যাপ্সার চিকিৎসার ওষধসমূহের উৎপাদনকারী প্রতিষ্ঠান আমদানী রেজিস্ট্রেশনের শর্তানুযায়ী উন্নত দেশসমূহn h_vt USA, UK, Japan, Germany, France, Australia and Switzerland-Gi Free Sale Certificate `llj Kii tZ cti bvb ejq qu j vBtmWys A_wi ll (Wmim) KZK.BZtcleqinRt-kb ctb vbi Kiv nq bvb/</i></p> <p><i>মানবিক কারণে ক্যাপ্সার আক্রান্ত- tivMf`i mijf Pukrmvi `#_etkI wetePbvq GB mKj Ati Ril÷KZ.c`mgf ctk®Zp t`k eñZ Ab`v t`k nBtZ ikij kZAS-eZtRyj xb Avg`vbi Avte`b wetePbv Rb~ we~vñZ Avtj vPbv ceR GB gtg®mgvñi k গৃহীত হয় যে, পশ্চ চিকিৎসার ওষধের ক্ষেত্রে বর্তমান প্রচলিত রেজিস্ট্রেশন প্রতিক্রিয়া অনুসরণ পূর্বক কেবলমাত্র তালিকাভূত 24 (PteYk) ll DbiZ t`tki gta~ th tKib 01 (GK) ll Drm t`tki wd*tnj mñJdtkUi wñEtZ weia tgvZtkeK Ab`v b~ kZtjej x ctY Kii qv wetc kI wetePbvq Avg`vbi tIRt-kbti mthwM ctb vbi Kiv hvBtZ cti   KivYvi Pukrmvi JIai mijf gj~ I ctc~Zv ibioZKij Yit~GB ikij রেজিস্ট্রেশন প্রতিক্রিয়া কেবলমাত্র পরবর্তী ০৬ (ছয়) মাস সময়সীমা পর্যন্ত বলুক থাকিবে।</i></p> <p><b><u>24 ll t`tki Zwi Kiv ibqSY t</u></b></p> <p><i>(1) At~ij qv (2) Añ~tq (3) tej wRqg (4) KvbWw (5) tWbgyK®(6) wdbj vU (7) dñY (8) Rvgfbx (9) nvt½ix (10) Avqij vU (11) BZyj x (12) Rvcib (13) tb~vij vU (14) bi l tq (15) mñarcjy (16) mñtWb (17) mñRvij vU (18) hñivo (19) hñiR (20) iñkqy (21) tcyj vU (22) t`ub (23) tPK wñcveij K Ges (24) `ññY tKwii qv/</i></p> <p><i>tgmwm®ZRviZ tnj &amp; tKqvi gvtKls Gi Avte`b wetePbv Rb~ Dc~vcb Kiv nq/</i></p>	<p><i>RvZiq JIa bñZ-2005 Gi mt½ cñ~vbi msMñZcY®bq weavq Avte`b bvgÄj Kiv hvBtZ cti  </i></p>	<p><i>RvZiq JIa bñZ-2005 Gi mt½ cñ~vbi msMñZcY®bq weavq Avte`b bvgÄj Kiv nBj  </i></p>

bs	eeiY	tUKibKij mve-Kigui 56 Zg mfvi mxvš-	mfvi mxvš-
2.13 /	<p><i>mvtbwid GtfwUm eisj vt` k ij t fikimib (Bbdzqbr fikimb) fwm tbi</i> Strength Ges      Indication <i>AcwieZ _mkij iaystrain cweZbi KvitY bZb c` mmite metePbri br Kwi qv      tivRt÷kb c0vtb c0vle Kwi qvtQ /</i></p> <p>তাহারা পত্রে উল্লেখ করিয়াছে যে, প্রতি বৎসর WHO KZR.mvi cwexi Rb' wzb Strain Gi Bbdzqbr fikimb wbePb Kiv nq/ wbePb Strain ,ij nBtZtQ H1N1, H3N2 Ges Type-B / D<sup>3</sup> Strain mgv mutagenic KvitY cweZ nq /</p> <p><i>c0Z0rbwU wgenwZ Composition Gi Bbdzqbr fikimb 238 Zg DCC-Gi mfviq Abfgv'b      cwBqv'tQ /</i></p> <p>A/ Brisben /59/2007 (H1N1)      A/ Brisben /10/2007 (H3N2)      B/ Florida/ 4/ 2006</p> <p>তাহারা পত্রে উল্লেখ করিয়াছে যে, উ<sup>3</sup> fikimb 2009-2010 Gi Rb' WHO wgenwZ Composition c0vle Kwi qvtQ hwn EMEA KZR Abfgw' Z /</p> <p>A/ Brisben /59/2007 (H1N1)-Like Strain      A/ Brisben /10/2007 (H3N2) -Like Strain      B/ Brisben /60/2008 -Like Virus</p> <p><i>fwm b-Gi Strength Ges Indication AcwieZ _mkij iaystrain cweZbi Rb' DCC Gi      Abfgv'b c0vRb nBte wkbri, mxvtsli Rb' Dc' tcb Kiv nq /</i></p>	<p><i>GB RvZiq fKimbti Avte'b      metePbri Rb' WMM KtUij KiguiUtz      Dc' tcb Kiv hvBtZ c0vle /</i></p>	<p><i>GB RvZiq fKimbti Avte'b      metePbri Rb' WMM KtUij      KiguiUtz Dc' tcb Kwi tZ nBte /</i></p>

<i>bs</i>	<i>meeiY</i>	<i>tUKibK'ij mve-KigilUi 56 Zg mfvi m̄m̄xvšl</i>	<i>mfvi m̄m̄xvš-</i>
2.14/	<p>M/s. Beximco Pharmaceuticals Ltd. <b>KZK Aitev Z Tadalafil 40 mg Tablet</b> (Tadalafil INN 40 mg Ges M/s. Novo Healthcare Ltd. <b>KZK Aitev Z Sildenafil 25 mg Tablet</b> (Sildenafil Citrate INN 35.119 mg eq. to 25 mg Sildenafil) Ges <b>Sildenafil 50 mg Tablet</b> (Sildenafil Citrate INN 70.238 mg eq. to 50 mg Sildenafil) <b>Abtgr`tb i weItq MZ WMM K̄bUj KigilUi 238 Zg m̄fīq Dc `vcb Kiv nBj - JI aui Safety, Efficacy and usefulness m̄utK®e - m̄i Z Z_ I DcIEmn JI a ibqSY KigilUi tUKibK'ij mve-KigilUi eivei c̄z̄te`b `wl̄tj i Rb` ibgevVZ m̄tklĀt i mḡtq GKU KigilU MVb Kiv nqt</b></p> <p>(1) Aa`vcK tgvt Bmgibj Lv, d̄igR Kij R efM, XvKv tgM̄tKj Ktj R - AvneqK  (2) Aa`vcK d̄ii`v teMg, tUKibK'ij d̄igR Kij R efM, XvKv wekje`ij q - m`m"  (3) Rbve Ate`jy ḡyW i, gnv-mPe, eisj vt`k JI a ik̄i m̄giZ Ges e`e`vcv c̄i Pij K, Bb̄tmPv d̄igRmD tUK'ij m̄w̄j t - m`m"  (4) Aa`vcK tMjv ḡKem qv, BDtivj Rx efM, eheUztkL ḡRe tgM̄tKj wekje`ij q/  (5) Rbve tgvt i "uj Avigb, JI a Z̄yeaqK, JI a c̄kmb c̄i`Bi, XvKv - m`m"-mPe/</p> <p>D<sup>3</sup> KigilU ibgiēc gZigZ c̄vib Kii qitQ t</p> <p>Though it is a scientific drug which is a PDE inhibitor which helps in erectile dysfunction, still we don't recommend it for registration in Bangladesh. The reason is chance of misuse/abuse in our country as there is no restriction facilities for prescribing by authorized prisciber in our country. If it is not properly advised by appropriate specialist for appropriate medication that may create social problem like other narcotic drugs (addictive drugs).</p>	<p>Ace`envi nBtZ c̄ti weaq m̄ḡiRK tcl̄iCU weTePbv Kii qv Ate`b tMZ ivLv h̄BtZ c̄ti /</p>	<p>Ace`envi nBtZ c̄ti weaq m̄ḡiRK tcl̄iCU weTePbv Kii qv Ate`b tMZ ivLv nBj /</p>

3/ AiZ DcKvix cÖYx kKb i¶v Ges Aib\_ t\_ mn Abib\_ tivtMi cÖKvc eÜ Kwi evi j+¶ evsj vt`tk WbtkdbvK Drcv`b eÜ Kwi evi Ri " ix c+¶c MÖY cÖt½/

“” I ciw evi Kj “gšyij tqi “gi K bs-Rb “”-1/Jla-6/2007/281, ZwiL t 03-10-2010-G (cZikv ÖM) Gi gva tg miPe, ciw tek I eb gšyij q nbtZ cÖB 15-09-2010 ZwiLi 464 msL’K wvi cÖU (cZikv ÖM) Jla ckvmb Awa`Bti tcöY Kiv nq/ D3 পত্রে উল্লেখ করা হয়েছে যে, বিজ্ঞানীদের মতে শকুন কমছে ব্য j qvB ci i Aib\_ , h2v, Rj vZ½, Ljv tivM ইত্যাদি বৃদ্ধি পাইতেছে। ইহা হইতে মানুষও আক্রান্ত হইতেছে। আমাদের দেশে “” bZvi cte cÖq 50,000 kKb nq/ GLb gvI 2000/ Zvi Averi cÖZv`b KigtZtQ/ iKb Kj GB gnv cÖtqi cÖvb Kvi Y ci PwKrmvq e\_vbvkK ØWbtkdbvK Gi eenvi / gwKØ MteIK W. wj i tm IK 2003 mvtj cÖvb Ktib th WbtkdbvK Jla eenvi Kiv nBqitQ Ggb Mi “” Qmjtj i gZt`n f¶Y Kwi t j kKtbi wKbx bó nBqv AwPtiB kKb gvi hvq/ tm Kvi t Y 2006 mvtj fviZ I cÖKvib Ges 2009 mvtj tbciy WbtkdbvK eÜ Kvi qv Gi ciw teZÖtqj wv KvgØ eenvi i m×vSjwbqitQ/

mi Kvit i m×vSj Avtj vtK AiZ DcKvix cÖYx kKb i ¶v\_ Jla ckvmb Awa`Bti i “gi K bs-WG/mewa-3/03/11273, ZwiL t 11-11-2010 (cZikv 000) tgvZiteK BmjKZ. Awdm Avt`k Øvi ØWbtkdbvK0 RvZxq tfUvibvix Jla Drcv`b NMZ Kiv nq/ D3 Avt`tk BtZvgta” Drcw Z Jlamgn Awmgx 27-04-2011 ZwiLi ci evRvi RvZ Kiv hvBte bv ejj qv wbt`R t`l qv nq/

“” I ciw evi Kj “gšyij tqi “gi K bs-Rb “”-1/Jla-6/2007/329, ZwiL 24-11-2010 (cZikv ÖPØ) ctí উল্লেখ করা হয়েছে যে, সরকার এই মর্মে সিদ্ধান্ত গ্রহণ করিয়াছে যে, “শকুন ও জনস্বাস্থ্যের Rb” ¶ZKi tfUvibvix Jla WbtkdbvK tmwWqvg Drcv`b eÜ Kiv nBte Ges BtZvgta” Drcw Z WbtkdbvK tmwWqvgi weecbb 6 gvm ci eÜ Kiv nBteØ/ gšy mfvj eVtK MpxZ D3 tfUvibvix WbtkdbvK tmwWqvg etÜi m×vSj ev`evqtbj Rb” Rj ix c+¶c MÖY Kiv `i Kvi / D3 পত্রে আরোও উল্লেখ করা হয় যে, The Drug Control Ordinance 1982 Ablyqri j vBtmWys A\_wiU Jla wqSY Kigui mgwi tk i Avtj vtK th tKb JItai tiRt÷kb ewZj Kwi tZ ci i / tmB Avtj vtK AvBbMZ wKj ¶ iwlqj D3 Kigui mgwi Avnevb Kvi qv mi Kvit i m×vSj ev`evqtbj c+¶c MÖY Kwi evi Rb” gnvcwi Pj K, Jla ckvmb Awa`Bi tK Abjvva Kiv nq/

GgZve “”q, evsj vt`tk Drcw Z I evRvi RvZKZ. ci PwKrmvq e“eüZ WbtkdbvK tmwWqvg RvZxq JItai tiRt÷kb ewZtj i we l tq m×vSjMÖtYi Rb” Dc“lcb Kiv nq/

mfvj Dcwi D3 we l tq we “”i Z Avtj vPbv nq/ ডাইক্লোফেনাক সোডিয়াম জাতীয় ঔষধের পার্শ্বপ্রতিক্রিয়ায় শকুনের কিডনী bó nBqv hvq weavq WbtkdbvK tmwWqvg RvZxq tfUvibvix JItai tiRt÷kb ewZtj i ct¶ m`m MY gZ cÖKv Ktib/

m×vSj t ci PwKrmvq e“eüZ mKj cÖvi tWtmm dtgØ WbtkdbvK RvZxq JItai tiRt÷kb ewZtj i mgwi k Kiv nq/ GB Avt`k 27-04-2011 ZwiL nBtZ KvhRi nBte/

4/ meva Avtj vPbv I m×vSj t

K) e½eÜztkL gJRe tgWtKj wekje`vj tqi DcPhW. cÖY tMvciy `E mfvj Ktib th, t`tk tek wKoY cÖövb wDg“vb I fvtUvibvix Jla Drcv`b Kvi qv \_vK/ D3 cÖvi Jla Aek`B c„K e“e“lq Drcv`b Kiv Avek`K/ cÖövbmgñ KZR. wDg“vb I fvtUvibvix RvZxq Jla, wj h\_vh\_ c„K e“e“lq Drcv`b Kiv nq wKb Zvw gj“qY ceR cÖZte`b `wLtj i Rb” wbtgæwYZ wekti ÁMtYi mgŠtq GKU Kigui MVb Kiv nq t

- (1) Aa“vck Wv. tgvt BmgvBj LvB, dvgtKv j R wefim, XvKv tgWtKv j Ktj R, XvKv/
- (2) Aa“vck dwi vteMg, wKtBv j dvtgmx I dvgtKv j R wefim, XvKv wekje`vj q/
- (3) cÖZvba, cÖYx m¤u` Awa`Bi, XvKv/
- (4) Wv. tgvtgbjv nK, wmbqj mn mfvcwZ, evsj vt`k Jla wki mgvZ/

L) *mfvq Aa'icK Wl. tgvt BmgvBj Lvb ej b th, JIa mbqSY KigUJi 236 Zg mfvq GB gtg@m×vsl i inqvtQ th, t`tk ~vbxq Drcv`tbi Rb" bZb JIa Ges cØvij Z JI tai bZb gvvi Avte`bKZ.c`mgtni gj~vqtbi t¶tî আবেদনকারীগণ কর্তৃক সংশিগ্ন পদসমূহ USFDA KZR Abtgr`b A\_ev BNF-G Aš-fP A\_ev t`tk cØqvRb i inqvtQ GB aitbi JI tai t¶tî Dchj cØqvbw/Z\_mn Avte`b Kwi‡Z nBte/ Safety, usefulness, efficacy-Gi Avtj vtK gj~vqb Kwi‡Z Avte`bKZ.c`mgtni USFDA KZR Abtgr`b A\_ev BNF-G Aš-fP A\_ev t`tk cØqvRbxq i inqvtQ GB cØvij JI tai t¶tî Dchj cØqvRb Z\_mn/tiditYi vfiE‡Z tUKibK'ij mve-KigUJiZ Dc ~vcb Kwi‡Z nBte/*

*vZib ej b USFDA GKU mbqSYKvix KZ@P Abv`tK BNF GKU PDR (Physician Desk Reference), vZib BNF Gi cweitz©UK-Gi JIa mbqSYKvix KZ@P MHRA (Medicines and Healthcare products Regulatory Agency) Gi Abtgr`tbi Z\_ `vLti ct¶ gZ cØvk Ktib/ mfvciZ gtni`q JIa mbqSY KigUJi tUKibK'ij mve-KigUJi mfvq D³ weI tq we ~vbi Z Avtj vPbv Kwiqv gZvgZ cØvibi Rb" ej v nBj/*

*m×vsl t BNF Gi cweitz©UK-Gi JIa mbqSYKvix KZ@P MHRA (Medicines and Healthcare products Regulatory Agency) KZR Abtgr`tbi Z\_ `vLti weI tq tUKibK'ij mve-KigUJi mfvq we ~vbi Z Avtj vPbv Kwiqv gZvgZ cØvibi Rb" ej v nBj/*

*tgRi tRbuij tgvt Avej Kyj ug AvRv`  
gnvcwi Pvj K  
JIa cØvmb AwA`Bi  
I  
m`m"-mip  
JIa mbqSY KigUJi/*

*gyas ûgvqly Kri  
mip  
-r- I cweivi Kj iiY gSYij q  
I  
mfvciZ  
JIa mbqSY KigUJi*